

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Conqueror[®] Herbicide

ACTIVE CONSTITUENTS: 300 g/L TRICLOPYR present as the butoxyethyl ester
100 g/L PICLORAM present as hexyloxypropylamine salt

GROUP	I	HERBICIDE
--------------	----------	------------------

For the control of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use Table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT
SHAKE WELL BEFORE USE.**

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road, Laverton North Victoria 3026
Tel: (03) 9282 1000
nufarm.com.au

APVMA Approval No.: 57783/108132

Contents: 1L, 5L, 10L, 20L, 110L, 1000L

STORAGE AND DISPOSAL (5L, 10L, 20L)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable containers (1000L only)

Store in the closed, original container in a cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

May cause an allergic skin reaction. Causes serious eye irritation. Combustible liquid. Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective eye and face protection. Avoid breathing spray. Contaminated work clothing should not be allowed out of the workplace. If eye irritation occurs and persists: Get medical advice/attention. IF SWALLOWED: Rinse mouth. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention.

In case of emergency: Phone 1800 033 498 Ask for shift supervisor. Toll free 24 hours.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or from the Nufarm website – nufarm.com.au

Conditions of sale

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS - AUSTRALIA WIDE
---	--

Conqueror® trade mark owned or used under license by Nufarm Australia Ltd.

BN / DOM / Barcode

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Conqueror[®] Herbicide

ACTIVE CONSTITUENTS: 300 g/L TRICLOPYR present as the butoxyethyl ester
100 g/L PICLORAM present as hexyloxypropylamine salt

GROUP I HERBICIDE

For the control of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use Table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT
SHAKE WELL BEFORE USE.**

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road, Laverton North Victoria 3026
Tel: (03) 9282 1000
nufarm.com.au

APVMA Approval No.: 57783/108132

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.

DO NOT spray if rain is likely within one hour or if foliage is wet from rain or dew.

DO NOT burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.

DO NOT apply by aerial application in wind in excess of 15km/hr and air temperatures above 35°C. In areas prone to flooding, treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.

1. WOODY WEED SITUATIONS

Table A: High Volume Spraying

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
African boxthorn	Less than 2m tall	All States	500mL/100L	Apply when bushes have good leaf cover, growth and no leaf fall.
<i>Angophora</i> spp.	1-3m tall		350mL/100L	
Australian blackthorn	Less than 2m tall		500mL/100L	Apply from late spring to early autumn.
Banksia spp.	1-3m tall		350mL/100L	
Biddy bush (Chinese shrub) (Sifton bush)	Autumn when actively growing	ACT and NSW only	500mL/100L	Add BS 1000 at 125mL/100L of water for best results.
Blackberry in association with: Docks, Ragwort, Smartweed Thistles	Late spring to autumn	All States	350mL or 500mL/100L	Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry. Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Conqueror.
Blue heliotrope	Flowering	NSW and Qld only	500mL/100L	Apply in a minimum spray volume of 1250L/ha
Brooms: Cape, English, Flax leaf, Montpellier	Spring to mid-summer prior to pod formation	All States	250mL/100L	Apply as a thorough foliage spray
	Autumn to winter		350mL/100L	
Camphor laurel	Less than 2m tall	All States	500mL/100L	
	Above 2m tall		350mL/100L	
<i>Casuarina</i> spp.	1-3m tall			
Chinee apple	Less than 2m tall	Qld and WA only	350mL/100L	Add BS 1000 at 100mL/100L of water for best results.
Cockspur thorn	Spring to Autumn	ACT, NSW and Qld only	350mL/100L	Apply as thorough foliage spray
Common sensitive plant	Any time when actively growing	NT, Qld and WA only	200mL/100L	To avoid leaves closing during application, spray plants while moving forward. Add BS 1000 at 100mL/100L of water for best results.
Crofton weed	Spring to autumn	ACT, NSW and Qld only	350mL/100L	Apply as thorough foliage spray.
Eastern cotton bush (<i>Maireana microphylla</i>)	Spring to autumn	NSW and Qld only	500mL/100L	Add Supercharge Elite at 500mL/100L of water. Some bushes may require a follow up spray to control regrowth.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Eucalyptus spp.	Seedling and regrowth from small lignotubers, 1-3m tall	All States	350mL or 500mL/100L	Apply the high rate where difficult to control species of Eucalyptus regrowth is present. Addition of an adjuvant may improve results - contact Nufarm Australia Limited for details.
Galenia	Fresh growth in spring to summer	NSW only	500mL/100L	Use 200L of spray mixture/ha
Giant bramble	Spring to autumn	NT, Qld and WA only		Penetration of thick clumps may be difficult and respraying may be necessary. Add BS 1000 at 100mL/100L of water for best results.
Gorse	1-1.5m tall	All States	250mL/100L	Spring and Summer treatment only. Add BS 1000 at 100mL/100L of water for best results.
	Over 1.5m tall or autumn treatment		350mL/100L	
	Winter treatment		500mL/100L	Brownout may not be complete until summer. Add BS 1000 at 100mL/100L of water for best results.
Groundsel bush (<i>Baccharis halimifolia</i>)	1-1.5m tall in spring to summer	All States	250mL/100L	Apply as a thorough foliar spray.
	Over 1.5m tall or autumn treatment		350mL/100L	
Green cestrum	Late spring to early autumn	ACT, NSW and Qld only	500mL/100L	One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high.
Hawthorn	Less than 2m tall	All States		Apply from late spring to early autumn.
Horehound	Pre- flowering	NSW and Qld only	350mL/100L	Apply as a thorough foliar spray
Japanese sunflower				
Lantana (<i>Lantana camara</i>) (<i>Lantana montevidensis</i>)	Up to 1m tall in summer to autumn	All States	500 or 750mL/100L	Add one of the following adjuvants, when using 350mL rate: A Supercharge Elite @ 0.5% v/v. Pulse® Penetrant @ 0.1% v/v. Thoroughly wet foliage, stems and soil around the base of the plants. Use higher rate on known harder to kill varieties.
	1-2 m tall in summer to autumn			
Lion Tail (<i>Leonotis nepetifolia</i>)	Pre-flowering	Qld only	200mL/100L	Apply as a thorough foliar spray. Add BS 1000 at 100mL/100L of water for best results.
Limebush	Any time of the year with good leaf cover and soil moisture	NT and Qld only	350mL/100L	Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results - contact Nufarm Australia Limited for details.
Manuka	At flowering	Vic only	500mL/100L	For optimum results, add Pulse Penetrant at 200mL/100 L of spray. Thoroughly wet foliage, stems and soil around the base of the plants.
Mesquite (<i>Prosopis</i> spp.) <i>Prosopis velutina</i>	Seedling, full leaf and flowering before podding	NSW, NT, Qld and WA only	350mL/100L	DO NOT spray plants bearing pods. Add BS 1000 at 100mL/100L of water for best results.
		Qld only	670mL/100L	
Mistflower	Spring to autumn	ACT, NSW and Qld only	350mL/100L	Apply as a thorough foliar spray.
Mother-of-millions	Flowering	NSW and Qld only	500mL/100L	Add BS 1000 at 100mL/100L of water
Paddy's lucerne	Active growth	NSW only		Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.
Parkinsonia	Under 2m tall	NT, Qld and WA only	350mL/100L	Add Supercharge Elite Oil at 500mL/100L water. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.
Prickly pear (common) Smooth tree pear	Active phyllode growth	All States	500mL/100L	Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.
Rubber vine (Not infected with rust)	Up to 1.5m tall at flowering	NT, Qld and WA only	350mL/100L	Spray all leaves and stems just to the point of runoff and thoroughly spray the base of the plant. With larger, more dense stands regrowth may appear. Subsequent control of any regrowth should be done by basal bark spraying.
	Dense stands greater than 1.5m tall at flowering		500mL/100L	
Siam weed	Active growth	Qld and WA only	350mL/100L	Add BS 1000 at 100mL/100 L of water for best results.
Sicklepod	Up to flowering	NT, Qld and WA only	200mL/100L	DO NOT apply to podding plants. Add BS1000 at 100mL/100L of water for best results.
St. John's wort	From flowering to early seed set	All States	500mL/100L	Late spring to early summer.
Sweet briar	Up to 1.5m tall	All States	350mL/100L	Add Associate® at 10g/100L water to obtain more reliable results with the lower rate of Conqueror. Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
			500mL/100L	
Tobacco weed	Actively growing plants	NT, Qld and WA only	300mL/100L	Add BS 1000 at 100mL/100L of water for best results.
Wattle (<i>Acacia</i> spp.) (except corkwood wattle)	1-3m tall	All States	350mL/100L	Add BS 1000 at 100mL/100L of water for best results.
Wild rosemary (<i>Cassinia laevis</i>)	Active growth, 0.5-1m tall	Qld only	350-500mL/100L	Use lower rate on seedlings 0.5m tall. Apply as a thorough foliar spray.
Wild tobacco tree	Spring to autumn up to 2m tall	ACT, NSW and Qld only	350mL/100L	

Table B: Aerial Application

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ha	CRITICAL COMMENTS
Blackberry	Summer to autumn	NSW, Qld, SA, Vic and WA only	10L	Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before re spraying with Conqueror. WARNING: <i>Eucalyptus</i> species up to 4m may be killed if sprayed during this treatment. Mature trees which are 15-20m tall may be partially defoliated but are likely to recover.
Gorse		Tas only		Helicopter application only
Cockspur thorn, Crofton weed, Lantana, Mistflower	Late autumn	NSW, NT and Qld only (helicopter only)	1.5L plus 6L Amine 625	Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150L/ha. Follow-up respraying will be required.
Lantana			10L	Helicopter application only
Rubber vine (Not infected with rust)	When flowering	NT and Qld only (helicopter only)	3-5L	Use rates will depend upon the density and height of the rubber vine stand. The higher rate should be used on dense stands. However, complete coverage and penetration may be difficult. Follow-up respraying will be required. Any regrowth should be sprayed with a suitable basal bark herbicide.
St. John's wort	Flowering to early seed set (Nov-Jan)	NSW only	4L	Helicopter application only. Follow up spraying will be required in the following season.
AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Parkinsonia	Seedlings 1-2m tall, or 12-24 months old	Qld and NT only (helicopter only)	3L	Add Supercharge Elite at 1 L/ha

Table C: Controlled Droplet Application (C.D.A.)

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Blackberry in association with: Docks, Ragwort, St. John's wort, Thistles	Summer to autumn	All States	Apply undiluted	One application may give satisfactory control but subsequent regrowth and seedlings should be resprayed after hardening off. Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Conqueror.

Table D: Low Volume High Concentrate Application Techniques (GAS POWERED GUN, SPRINKLER SPRAYER)

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Blackberry	Late Spring to autumn	ACT, NSW, Qld, SA, TAS and WA only	335mL/10L	Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
Camphor laurel, Cockspur thorn, Crofton weed	Less than 1.5m high	ACT, NSW and Qld only	500mL/10L	
<i>Eucalyptus</i> species	Seedlings up to 2m tall	All States	335mL/10L	Gas powered Gun only: Apply to actively growing bushes not more than 1.5m tall that have not more than 5 stems from the crown.
Mistflower	Less than 1.5m high	ACT, NSW and Qld only	500mL/10L	
Sweet briar	1.5m tall, full leaf to ripe fruit	NSW only		
St. John's wort	During flowering to early seed set	NSW, Tas and Vic only		Gas powered Gun only: One application should provide control. Minor regrowth and seedlings may be retreated the following summer.
Wild tobacco tree	Less than 1.5m high	ACT, NSW and Qld only	500mL/10L	Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.

Table E: Boom Application

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Galenia	Fresh growth during spring to summer	NSW only	5L/ha	Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
Sicklepod	Up to flowering	NT and Qld only	3L/ha	DO NOT apply to podding plants. Add BS 1000 at 100mL/100L of water.
St. John's wort	Flowering to early seed set (Nov-Jan)	NSW only	2-4L/ha	Use the higher rate on dense infestations and when longer residual control is required. Follow-up respraying will be required in the following season.

2. FALLOW SITUATIONS

Table F: Boom Application

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Blackberry nightshade-Suppression only	10-25cm tall prior to flowering	NSW and Qld only	200-400mL/ha + 1.2L Gladiator + Collide® 700 or 1L Gladiator OptiMax	FOR USE BY GROUND EQUIPMENT ONLY. Plants must be actively growing. Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5 cm diameter for <i>Polymeria pusilla</i>). Refer to mixture product label for use of Collide® 700 Surfactant. DO NOT plant susceptible crops for up to nine months after application, as specified in General Instructions - Minimum Recropping Periods - Black Cracking Clay Soils, Northern NSW & Qld. Dry conditions after application will increase the recropping interval.
Camel melon, Prickly paddy melon, Cucumber melon (<i>Cucumis melo</i>)	From 2 leaf to 50cm diameter			
Common sowthistle	From 8 leaf to flowering			
Cow vine	From 2 to 5 leaf up to 15cm diameter, prior to flowering		300- 500mL/ha + 1.2L Gladiator + Collide® 700 or 1L Gladiator OptiMax	
Lucerne (established)	Active growth, 15 to 25cm high, during spring			
<i>Polymeria pusilla</i>	2 to 12 leaf up to 20cm diameter, prior to flowering			

Table G: Blanket Wiper Application

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Bitterbark (<i>Alstonia constricta</i>)	From summer to end of autumn	Qld	1:4 (1 part Conqueror to 4 parts water) 2% solution for spot spray (e.g. 100mL Conqueror in 5L water)	For use with blanket wipers only. For best results apply in autumn to tall (>60 cm) plants using two opposite directional passes (up and back). Follow up "missed" plants with a spot spray application. These will be obvious after 6 weeks. Blanket wiper application can be made in summer when plants are smaller but follow up spot spraying may be necessary. Do not disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months). Spot spraying "missed" plants: thoroughly wet all stems and leaves without producing any solution run off. Avoid any spray reaching the soil surface.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

IN TASMANIA FOR BLACKBERRY: DO NOT treat bushes carrying mature or near mature fruit.

FOR NATIVE VEGETATION: Use of Conqueror on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

WITHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

FALLOW SITUATIONS

MINIMUM RECROPPING PERIODS - Black Cracking Clay Soils, Northern NSW & Qld.

Table F: Boom Application

Plant-back periods for crops following the application of Conqueror for rates up to 600mL/ha				
RATE mL/ha	200	300	400	600
CROP	MONTHS			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba Beans	4	4	6	6
Chickpeas	4	6	6	6
Lucerne	6	9	9	9

These plant back periods are based on a normal rainfall pattern. During drought conditions (or when the rainfall is less than 100mm for a period of 4 months or greater) the plant back period may be significantly longer.

Table G: Blanket Wiper Application

Plant-back periods for crops following blanket wiper application.	
CROP	MONTHS
Broadleaf crops	18
Lucerne	6
	This will allow any potential soil residues to dissipate, if any, and allow effective control.

NOTE: Before using Conqueror in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product with the longest plant-back period, will determine the time between spraying and planting.

COMPATIBILITY

Follow any regional restrictions, and all directions and restrictions on the label, for any chemical mixed with Conqueror (e.g 2,4,-D amine). Conqueror is compatible with the following herbicides: Amicide® Advance 700, Amine 625, Associate®, Gladiator®, Gladiator® OptiMax, weedmaster®ARGO®, weedmaster® DST®.

Conqueror is compatible with the following adjuvants, as per Directions for Use: BS1000, Collide® 700, Pulse® Penetrant, and Supercharge Elite.

MIXING

Mix only with water. Half fill the spray unit with water, and add the required amount of Conqueror. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

APPLICATION**1. WOODY WEED SITUATIONS**

Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.

Table A. High Volume Spraying

Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.

Hand Gun

Apply the recommended mix to give full coverage of leaves and stems through a No. 6-8 tip at 700-1500kPa (400-500kPa for St John's wort). A spray volume of 3000-4000L per infested hectare of 1-2 metre high blackberry (30-40L/100m²) should be used. Use 2000L of spray mixture/ha of galenia infestation (i.e. 20L/100m² infested area).

Knapsack

Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand gun. A spray volume of 3-4L/10m² infested area should be used. A spray volume of 2L/10m² should be used for an area infested with galenia.

Table B. Aerial Application

Apply in 200L of water/ha using an aircraft to apply 100L per pass on a double overlap pattern using nozzle configurations to produce droplets of 250-350 micron diameter. The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets. DO NOT spray when wind exceeds 15km/hr and /or air temperature reaches 35°C.

Table C. Controlled Droplet Application (C.D.A)

Results similar to high volume spraying can be obtained using Micron Herbi* or similar equipment. Select a nozzle to give a flow rate of 2mL/sec and sweeping action of approximately 1m/sec to ensure a droplet density of 20/cm². Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with C.D.A unit.

Table D. Low Volume High Concentrate Application Techniques

Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

Gas powered gun

Apply 50mL shots to obtain uniform coverage of 4-5m² of surface area of bush. This relates to 20 droplets/cm² of leaf surface.

Sprinkler sprayer

This technique involves using a micro sprinkler that is disconnected to a hollow fibre glass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50-200kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

Table E. Boom Application

Application in a minimum spray volume of 200L/ha for galenia and St. John's wort and 600L of water/ha for sickle pod. Flat fan nozzles are recommended, using pressure in the range of 200-300kPa. Boom height must be set to ensure double overlap of nozzle patterns.

2. FALLOW SITUATIONS**Table F. Boom Application**

Application of Conqueror in a minimum spray volume of 50L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200-300kPa. Boom height must be set to ensure double overlap of nozzle patterns.

Table G. Blanket Wiper Application

Blanket needs to be made from durable and wettable material with a rigid backing. Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (rewetting). Ideally, a scraper bar should be mounted at the front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping stems. This scraper may be mounted at the front of the vehicle. Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10-15 kph are ideal for blanket wiping application.



spraywisedecisions.com.au is an online weather forecasting program and is recommended for use when planning your pesticide application.

CLEANING SPRAY EQUIPMENT

Rinsing

After using Conqueror empty the spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles. After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsings procedure twice.

Decontamination

Before spraying cotton and other sensitive crops with equipment that has been used to apply Conqueror, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section. Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent such as Tank and Equipment Cleaner at 500mL/100L of water and circulate throughout the system for at least 15 minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

RESISTANT WEEDS WARNING

GROUP	I	HERBICIDE
-------	---	-----------

'Conqueror' Herbicide is a member of the pyridines group of herbicides. Conqueror has the disruptors of plant cell growth mode of action. For weed resistance management, Conqueror is a Group I herbicide. Some naturally-occurring weed biotypes resistant to Conqueror and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Conqueror or other Group I herbicides. Strategies to minimise the risk of herbicide resistance are available.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Crops susceptible to Conqueror include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and *Pinus* spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines. Conqueror is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment. Picloram, one of the active constituents in this product can remain active in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation. DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops. DO NOT apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants. DO NOT apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants. DO NOT move soil, which may have been treated to areas where desirable plants are to be grown.

PROTECTION OF LIVESTOCK

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down. Many plants remain poisonous after death, and stock should not be allowed access, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable containers (1000L only)

Store in the closed, original container in a cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If in eyes, hold eyes open, flood with water for at least 15minutes and see a doctor.

ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

May cause an allergic skin reaction. Causes serious eye irritation. Combustible liquid. Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective eye and face protection. Avoid breathing spray. Contaminated work clothing should not be allowed out of the workplace. If eye irritation occurs and persists: Get medical advice/attention. IF SWALLOWED: Rinse mouth. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or from the Nufarm website – nufarm.com.au

In case of emergency: Phone 1800 033 498 Ask for shift supervisor. Toll free 24 hours.

Conditions of sale

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

® All trade marks owned or used under license by Nufarm Australia Ltd.

* Other trade marks.