

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**




**Active Constituent:** 95 g/L AZOXYSTROBIN

For the control of various diseases of turf as per the Directions for Use

**GROUP K FUNGICIDE**

APVMA Approval No: 58324/1/0405 Pack size: 1 L  
APVMA Approval No: 58324/5/0805 Pack size: 5 L



UN NO. 3082 ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CONTAINS AZOXYSTROBIN) MARINE POLLUTANT, PACKING GROUP III, HAZCHEM 2X, CONTAIN SPILLAGE.

## GENERAL INSTRUCTIONS

### Fungicide Resistance Warning

HERITAGE MAXX Fungicide is a member of the strobilurin and related products group of fungicides. For fungicide resistance management the product is a Group K fungicide. Some naturally occurring individual fungi resistant to the product and other Group K fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group K fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Syngenta Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

### Application

Apply in a sufficient volume of water to achieve thorough coverage of all foliage. In turf this can generally be achieved with spray volumes of 4 to 8 L per 100 m<sup>2</sup>.

### Mixing

HERITAGE MAXX is a micro emulsion fungicide that mixes readily with water. Add the required amount of product directly to the spray tank and mix well.

### Compatibility

HERITAGE MAXX is compatible with Banner Maxx® and Daconil Weather Stik®.

### PRECAUTION

#### Re-entry Period

DO NOT enter treated areas until the spray has dried, unless wearing chemical resistant gloves and footwear.

#### PROTECTION OF LIVESTOCK

DO NOT graze treated turf/lawn, or feed turf/lawn clippings from any treated area to poultry or livestock.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

HIGHLY TOXIC TO AQUATIC LIFE. DO NOT contaminate dams, waterways or drains with the chemical or used containers.

DO NOT apply under meteorological conditions or from spraying equipment that could be expected to cause spray drift on adjacent areas, particularly wetlands, waterbodies or watercourses.

DO NOT apply when heavy rains or storms are expected within 4 hours.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

**Will irritate the eyes and skin. Avoid contact with eyes and skin.**

**When opening the container and preparing spray wear:**

- cotton overalls buttoned to the neck and wrist (or equivalent clothing),
- elbow-length PVC gloves and
- goggles.

**If applying by hand wear:**

- cotton overalls buttoned to the neck and wrist over normal clothing and
- elbow-length PVC gloves.

**Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.**

### FIRST AID

**If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.**

### MATERIAL SAFETY DATA SHEET:

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone 1800 067 108 or visit our website at [www.syngenta.com.au](http://www.syngenta.com.au)

### MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Syngenta has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

®Registered Trademark of a Syngenta Group Company

## DIRECTIONS FOR USE

### Restrictions:

DO NOT use in the home garden

DO NOT apply by aerial application methods

DO NOT use this product through any type of irrigation system or ultra-low volume spray system

Situation	Disease	Rate	Critical Comments
Turf	Helminthosporium Disease ( <i>Bipolaris</i> spp, <i>Drechslera</i> spp, <i>Exserohilum</i> spp), Grey Leaf Spot ( <i>Pyricularia grisea</i> ), Anthracnose ( <i>Colletotrichum graminicola</i> ), Red Thread ( <i>Laerisaria fuciformis</i> ), Fusarium Disease ( <i>Fusarium nivale</i> )	30 to 60 mL/ 100 m <sup>2</sup> applied in 4 to 8 L water	Apply in a sufficient volume of water to achieve thorough coverage of all foliage. Apply HERITAGE MAXX in a preventative fungicide program containing fungicides from different chemical groups. Begin applications before symptoms occur when conditions first favour disease and continue applications while conditions remain favourable for disease development. <b>Spray interval and rate</b> Apply no more than 2 consecutive HERITAGE MAXX applications at 14 to 21 day intervals for Helminthosporium and at 14 to 28 day intervals for other diseases. Use the shorter spray interval or apply the higher rate when climatic conditions remain favourable for disease development over prolonged periods. Use the shorter interval and the higher rate when conditions for disease development are optimum.
	Brown Patch ( <i>Rhizoctonia solani</i> )	60 mL/ 100 m <sup>2</sup> applied in 4 to 8 L water	<b>Resistance management</b> Disease control may be reduced if strains of pathogens less sensitive to HERITAGE MAXX develop. HERITAGE MAXX is best used in a preventative disease control program. Reliance on curative use of HERITAGE MAXX could lead to resistance. As a precaution, DO NOT apply HERITAGE MAXX for more than a total of 1/3 of the fungicide applications in one season. If consecutive applications of HERITAGE MAXX are used they must be followed by at least the same number of applications of fungicide(s) from a different group(s) before HERITAGE MAXX is used again in the current or following season.
	<i>Pythium</i> spp	60 mL/ 100 m <sup>2</sup> applied in 4 to 8 L water	Apply in a sufficient volume of water to achieve thorough coverage of all foliage. Apply HERITAGE MAXX in a preventative fungicide program containing fungicides from different chemical groups. Begin applications when conditions first favour disease development and before disease is present. Apply 2 consecutive applications at 14 to 21 day intervals. Use the shorter spray interval when climatic conditions remain favourable for disease development over prolonged periods. <b>Resistance management</b> Disease control may be reduced if strains of pathogens less sensitive to HERITAGE MAXX develop. HERITAGE MAXX is best used in a preventative disease control program. Reliance on curative use of HERITAGE MAXX could lead to resistance. As a precaution, DO NOT apply HERITAGE MAXX for more than a total of 1/3 of the fungicide applications in one season. If consecutive applications of HERITAGE MAXX are used they must be followed by at least the same number of applications of fungicide(s) from a different group(s) before HERITAGE MAXX is used again in the current or following season.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD: DO NOT GRAZE OR CUT FOR STOCK FOOD

### WARNING

HERITAGE MAXX is extremely phytotoxic to certain apple varieties.

DO NOT spray HERITAGE MAXX where spray drift may reach apple trees.

DO NOT spray when conditions favour drift beyond the area intended for application. Conditions that may contribute to drift include thermal inversions, excessive wind speed, certain sprayer nozzle/pressure combinations, small spray droplet size, etc.