

RAMPAGE®



APPROVED FOR FENCE LINE BAITING



PRODUCT FEATURES:

- ▶ Made with the powerful acute active ingredient Cholecalciferol (Vitamin D₃)
- ▶ Rodents cease feeding after consuming a toxic dose
- ▶ Substantially reduced risk of secondary poisoning
- ▶ Low toxicity to birds
- ▶ Excellent palatability
- ▶ Kills anticoagulant resistant rats and mice
- ▶ Weatherability
 - Stable up to 90° C
 - Mold and moisture resistant
 - Incorporates an advanced preservative system
- ▶ Pelleted Sachets
 - Ensure freshness and convenience
 - Must be used in bait stations
- ▶ Approved for perimeter fence line application

PRODUCT	PAIL QTY	CASE QTY
RAMPAGE® Rat & Mouse Bait	50 x 30g bait sachets 100 X 10g bait sachets	1.5 kg pail x 4 1 kg pail x 4



More Than Meets The Eye

Madison, Wisconsin 53704 USA | Ph: (608) 241-0202 | Fax: (608) 241-9631
australasian@belllabs.com

www.belllabs.com

RAMPAGE[®] 100 x 10 g Bait Sachets (1.0kg Pail)

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships), around grain terminals, storage areas and perimeter fence lines.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>)	6 to 18 bait sachets per bait station every 9 metres.	Eliminate, as far as practicable, all alternative food sources. Bait infested area. Place bait sachets in lockable tamper-resistant bait stations. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait sachets. Continue observation and replenishment until no more bait sachets are taken.
	Mice (<i>Mus musculus</i>)	3 bait sachets per bait station every 3 metres.	

RAMPAGE[®] 50 x 30 g Bait Sachets (1.5kg Pail)

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships), around grain terminals, storage areas and perimeter fence lines.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>)	2 to 6 bait sachets per bait station every 9 metres.	Eliminate, as far as practicable, all alternative food sources. Bait infested area. Place bait sachets in lockable bait stations. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait sachets. Continue observation and replenishment until no more bait sachets are taken.
	Mice (<i>Mus musculus</i>)	1 bait sachets per bait station every 3 metres.	

RAMPAGE[®]

RAT AND MOUSE BAIT

