



POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE
OPENING OR USING



International

BRIGAND

RODENTICIDE BLOCKS

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM

NET CONTENTS

8kg

READY TO USE BLOCK BAIT FOR USE IN DAMP OR DRY SITUATIONS
IN AND AROUND INDUSTRIAL, COMMERCIAL, AGRICULTURAL
AND DOMESTIC BUILDINGS TO CONTROL RATS AND MICE

CONTAINS
DENATONIUM
BENZOATE AS A
BITTERING AGENT

SINGLE FEED – CONTROLS SPECIES RESISTANT TO WARFARIN

Distributed in Australia by: **PelGar International (Aus) Pty Ltd.**

Northbank Plaza, Level 22, 69 Ann Street, Brisbane QLD 4000

Tel. 07 3112 1024 www.pelgar.co.uk

Manufactured by PelGar International Ltd.

BRIGAND RODENTICIDE BLOCKS

DIRECTIONS FOR USE

RESTRAINTS

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.
DO NOT place bait stations in the open.
DO NOT apply baits to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around Industrial, Commercial, Public Services, Agricultural and Domestic Buildings	RATS (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>)	1 to 3 blocks per bait station depending on level of infestation	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings.
	MICE (<i>Mus musculus</i>)		For Rats: Place 1 to 3 blocks at intervals of 5-9 m in infested areas. For Mice: Place single blocks at 2-3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.

GENERAL INSTRUCTIONS

Keep away from food and water, and surfaces that come into contact with food and water. Contains human taste deterrent, which prevents accidental consumption by children.

Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait the infested area. Place baits under cover. Inspect baits daily. Replace eaten baits. If quickly eaten, increase number of baits. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and locations of bait stations for purpose of replenishment and later removal. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased. The use of this product for the control of protected native rodents requires permission from wildlife authorities. The product is restricted to use in and around buildings (within 2 metres) or in enclosed spaces, e.g. drains.

Most rodent deaths occur 4 to 7 days after consumption of bait.

PRECAUTIONS

DO NOT feed the bait to wild or domestic birds.
DO NOT apply directly on or near exposed food or food utensils, food preparation areas, livestock feed or water, or food storage areas.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. Place baits in locations inaccessible to other animals or birds.
DO NOT contaminate dams, waterways and drains with bait or its used containers. Look for dead rats or mice – burn or bury them, do not place in refuse bins or rubbish tips.

STORAGE AND DISPOSAL:

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Dispose of rinsings and unused or untaken bait in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Ensure uneaten baits are removed and disposed of with empty containers when rodent activity has ceased.

SAFETY DIRECTIONS:

Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K1 (PHYTOMENADIONE) is antidotal.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone Australia 131126.

MSDS: Additional information is listed in the Material Safety Data Sheet available from the supplier.



® Brigand is a registered trademark of PelGar International Ltd.

APVMA APPROVAL NO. 68025/57306

Batch No. See bucket.

D.O.M. See bucket.

NET CONTENTS

8kg

