

Storm[®] Soft Bait

Rodenticide

A powerful, palatable and highly effective soft bait rodenticide

General description

Storm® Soft Bait Rodenticide has been developed into a unique, high oil content, soft bait formulation for unsuppressed efficacy even in the most difficult to control situations. Unlike other soft baits, Storm Soft Bait is formulated at the equivalent concentration as block formulations, making it the most potent soft bait formulation available.

The unique active ingredient in **Storm Soft Bait**, flocoumafen, is a second generation rodenticide with a high potency. The power of **Storm Soft Bait** means it only takes a single feed for both rats and mice to consume a lethal dose, giving fast and reliable control.

<text><text><image><section-header><section-header><section-header><section-header><text>

Storm Soft Bait is effective on both rats and mice and can be used in a variety of situations, including within and around domestic, commercial, agricultural and industrial buildings.

Features	Benefits			
Palatable formulation	Storm Soft Bait has been formulated into a unique, high oil content, soft bait formulation resulting in rapid bait uptake in the most difficult of environments.			
Rapid control	Storm Soft Bait has been formulated with the powerful active ingredient flocoumafen (0,05g/kg). Unlike other soft baits, this full strength loading results in rapid control, first time.			
Performance and reliability	The extremely palatable formulation is designed to improve the effectiveness in dry environments, and among rodent populations that have become reluctant to feed on traditional block baits.			

Application

Storm Soft Bait should only be placed in areas frequented by the target pests and within rodent burrows.

Storm Soft Bait can be used strategically with BASF's entire rodent control range, providing pest control operators and farmers with the most superior, unique and comprehensive, cleanout, rotation and resistance strategy available for all situations.

Packaging

Available in 5kg buckets.





Storm Soft Bait

Rodenticide

Situation	Pest	Rate	Critical comments
In and around industrial, commercial, agricultural domestic, and public service buildings, in sewers, animal houses, wharves and around grain terminals.	Rats - <i>Rattus rattus</i> also known as Black rat, Roof, rat, Ship rat or, Fruit rat and <i>Rattus norvegicus</i> also known as the Brown rat or Norway rat	3 - 5 Storm Soft Baits at each bait location 1 - 2 Storm Soft Baits in each rat hole/ burrow	 Select suitable bait locations, such as in active rat holes/burrows, along rat runs and under rubbish. Try to establish a barrier of bait locations between living and feeding areas. Secure 3 - 5 Storm Soft Baits at each bait location. Storm Soft Baits should be placed in areas inaccessible to non-target species, such as in bait trays, inside bait boxes or within enclosed spaces. 1 - 2 Storm Soft Baits can also be placed within each active rodent hole/ burrow. On initial application, an inspection of bait locations after 3 days may be undertaken to determine activity in bait locations and new locations may be selected if necessary. A subsequent inspection may also be undertaken again after 4 days to replace any soft baits which have been eaten. For severe infestations it is recommended that repeat inspections of all bait locations are conducted at 7 day intervals, replacing fresh Storm Soft Baits only when Storm Soft Baits have been eaten and until all signs of rat activity have disappeared. Normally only 3 to 4 baiting rounds (pulses) are required for control.
	Mice – <i>Mus</i> <i>musculus</i> also known as the House mouse	1 - 3 Storm Soft Baits at each bait location	Mice have very limited home ranges and do not need to drink. They are more difficult to control than rats because their feeding patterns are more erratic. Selection of bait locations is therefore even more important than for rats. Secure 1-3 Storm Soft Baits in areas inaccessible to non- target species, such as in bait trays, inside bait boxes or within enclosed spaces, spaced 2-3 metres apart. Bait locations should be in areas where activity is obvious, particularly where droppings are seen. Inspect bait locations and replace Storm Soft Baits as for rats until all signs of mouse activity have disappeared.



For technical advice and product information visit **www.pestcontrol.basf.com.au**