

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

 **Nufarm WEEDMASTER<sup>®</sup>**  
**DST<sup>®</sup>**

ACTIVE CONSTITUENT: 470 g/L GLYPHOSATE  
present as the potassium and mono-ammonium salts

**HERBICIDE**

**GROUP 9 HERBICIDE**

Water soluble herbicide for non-selective control of many annual and perennial weeds in conservation tillage and other situations.

**SL**

**SOLUBLE CONCENTRATE**



 **Nufarm**

**AUSTRALIAN  
THROUGH  
& THROUGH**

# INDEX

Section	Page Number
1. FALLOW AND PRE-SOWING – ALL CROPS	2
2. WEED TABLE FOR SECTION 1, SECTION 10 AND SECTION 13	2
3. FLEABANE CONTROL FOR FALLOW AND PRE-SOWING – MIXTURES WITH AMICIDE ADVANCE 700	12
4. CANOLA - HYBRIDS WITH THE OPTIMUM GLY® HERBICIDE TOLERANCE TRAIT	13
5. CANOLA - TRUFLEX* WITH ROUNDUP READY* TECHNOLOGY	14
6. CANOLA - ROUNDUP READY*	15
7. COTTON - ROUNDUP READY FLEX*	16
8. PRE AND POST HARVEST USES, PASTURE TOPPING	18
9. OPTICAL SPOT SPRAY TECHNOLOGIES	20
10. POST SOW PRE-EMERGENT (PSPE) USES, SHIELDED SPRAYERS AND BAND SPRAYING	20
11. BRUSH AND WOODY WEEDS – MIXTURES WITH ASSOCIATE	21
12. UNWANTED TREE CONTROL	21
13. GENERAL WEED CONTROL	22
14. WEED TABLE FOR SECTION 13 (HANDGUN, KNAPSACK ONLY)	23

## DIRECTIONS FOR USE

### RESTRAINTS

**DO NOT** use Nufarm weedmaster® DST® Herbicide, or any other product containing glyphosate, as the only method of weed control if glyphosate resistant weeds are suspected or present.

To ensure herbicide absorption, **DO NOT** disturb weeds by cultivation, sowing or grazing for 1 day after treatment of annual weeds and 7 days for perennial weeds, except where noted.

**DO NOT** treat weeds under poor growing conditions due to moisture stress, waterlogging, severe frosting, insect damage etc. Apply treatments to actively growing weeds which have at least one true leaf (broadleaf weeds) or two leaves (grasses) to provide an adequate surface area for herbicide uptake. Reduced performance may also occur where weeds are covered with dust or silt. If heavy grazing has occurred, allow regrowth to 6-8 cm before spraying and use the higher rates recommended (where higher rates are listed for the relevant crop/situation).

**CAUTION** Certain plants may be naturally toxic to stock. Where known toxic plants are present **DO NOT** allow stock to graze until complete browning of treated plants has occurred.

**DO NOT** apply by aircraft when temperature is above 30°C.

### SPRAY DRIFT RESTRAINTS

**DO NOT** apply by a vertical sprayer.

Specific definitions for terms used in this section of the label can be found at [www.apvma.gov.au/spraydrift](http://www.apvma.gov.au/spraydrift)

**DO NOT** allow bystanders to come into contact with the spray cloud.

**DO NOT** apply in a manner that may cause an **unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production**, or cause contamination of plant or livestock commodities, outside the application site from **spray drift**. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

**DO NOT** apply unless the **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application.

**DO NOT** apply if there are **hazardous surface temperature inversion** conditions present at the **application site** during the time of application. **Surface temperature inversion conditions** exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

### Section 1. FALLOW AND PRE-SOWING – ALL CROPS

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Prior to sowing any crop or pasture or for a fallow (refer to partner product labels for planting intervals)	Annual and perennial weeds including those listed in <b>Section 2 weed table</b> where minimum application rates are listed.	0.77-7.0 L/ha	Nufarm weedmaster DST may be used for control of annual and perennial weeds as directed, in agricultural land prior to sowing of any edible or non-edible crop (unless otherwise indicated on a tank mix partner label), except transplanted tomato seedlings. <b>DO NOT</b> apply prior to transplanting tomato seedlings.
		OR	On friable soils and where there is only light cover of young weeds, sowing may proceed satisfactorily from one day after spraying. In situations of heavy weed growth sowing should be delayed until weed decay and soil conditions allow formation of a satisfactory seed bed.
		0.77-2.3 L/ha + 0.28-1.2 L/ha Amicide® Advance 700	Nufarm weedmaster DST requires the addition of a tank-mix surfactant such as Activator or Collide 700 (refer to the Surfactant Addition section of the General Instructions), except where specifically noted in the Directions for Use. Rainfall occurring up to 6 hours after application may reduce effectiveness. Rainfastness or general efficacy may be reduced if weeds are not actively growing, are under stress or conditions of low light intensity/darkness. Heavy rainfall within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required. Delay treatment of plants wet with dew or rain, if water droplets run off when plants are disturbed. Avoid contact with foliage, green stems or fruit of crops, desirable plants and trees, since severe injury or destruction may result.
OR	0.77-2.3 L/ha + other tank mix partner products listed in Section 2 weed table (including Terrad® or® 700WG Herbicide)	Minimum application rates for each weed are listed in the table in Section 2 (in mL/ha); recommended use rates (except where a tank mixture is used where the maximum rate is 2.3 L/ha, or when applying by aircraft when the maximum rate is 3.1 L/ha) are: - 0.77-2.3 L/ha for annual weed species - 2.3-4.6 L/ha for perennial species - 7.0 L/ha for hard to control perennial species NOTE: The maximum approved rate for any weed is 7.0 L/ha and this should never be exceeded. For further information on application rate selection, please refer to General Comments following the weed table in Section 2 below.	

### Section 2. WEED TABLE FOR SECTION 1, SECTION 10 AND SECTION 13

Refer to the General Comments following this table prior to use.

WEEDS	MINIMUM RATE (mL/ha) <sup>A</sup>	CRITICAL COMMENTS
African lovegrass ( <i>Eragrostis curvula</i> ) #	2300	
African turnip weed ( <i>Sisymbrium thellungii</i> )	770	
Ageratum ( <i>Ageratum</i> species)	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 390 mL/ha <sup>B</sup>
Amaranth ( <i>Amaranthus viridis</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup>
Amaranth, dwarf (Amaranthus macrocarpus)	770	Always add Terrad® or® 700WG Herbicide at 15-40 g/ha with Banjo® Spray Adjuvant <sup>C</sup> at 1 L/100L to increase the speed of brownout and may also improve final control.
Amaranth, native ( <i>Amaranthus interruptus</i> )	770	Kamba® 750 at 105-160 mL/ha must always be added.

WEEDS	MINIMUM RATE (mL/ha) <sup>A</sup>	CRITICAL COMMENTS
Amaranth, redroot ( <i>Amaranthus retroflexus</i> )	2300	Alternatively to Nufarm weedmaster DST alone add either: - Terrad <sup>®</sup> or <sup>®</sup> 700WG Herbicide at 20-40 g/ha with Banjo <sup>®</sup> Spray Adjuvant <sup>^</sup> at 1 L/100L to increase the speed of brownout and may also improve final control - Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control.
Annual ground cherry ( <i>Physalis angulate</i> )	1530	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup>
Annual ryegrass ( <i>Lolium rigidum</i> ) #	1150	Alternatively to Nufarm weedmaster DST alone add Terrad <sup>®</sup> or <sup>®</sup> 700WG Herbicide at 40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST for higher levels of final control of annual ryegrass including populations with weak levels of glyphosate resistance.
Apple of Peru ( <i>Nicandra physalodes</i> )	1400	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup> - Kamba 750 at 105-160 mL/ha for improved control.
Artichoke thistle ( <i>Cynara cardunculus</i> )	2300	
Australian bindweed ( <i>Convolvulus angustissimus</i> ) *	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 280 mL/ha <sup>B</sup>
Australian bluebell ( <i>Wahlenbergia stricta</i> )	770	
Australian crassula/stonecrop ( <i>Crassula spp.</i> )	2300	Nail <sup>®</sup> 600EC Herbicide must always be added at 10-30 mL/ha.
Awless Barnyard grass ( <i>Echinochloa colona</i> )	770	Under Summer (hot) conditions, dense infestations may require follow-up treatment for complete control.
Ball mustard ( <i>Neslia paniculata</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 280 mL/ha <sup>B</sup>
Barbed wire grass ( <i>Cymbopogon refractus</i> )	2300	
Barley grasses ( <i>Hordeum</i> species)	770	Alternatively to Nufarm weedmaster DST alone add Terrad <sup>®</sup> or <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo <sup>®</sup> Spray Adjuvant <sup>^</sup> at 1 L/100L to increase the speed of brownout and may also improve final control.
Barnyard grass ( <i>Echinochloa crus-galli</i> ) #	1530	Under Summer (hot) conditions, dense infestations may require follow-up treatment for complete control. <b>DO NOT</b> tank mix with atrazine when spraying. Alternatively to Nufarm weedmaster DST alone add Terrad <sup>®</sup> or <sup>®</sup> 700WG Herbicide at 40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control.
Bathurst burr ( <i>Xanthium spinosum</i> )	1150	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Bellvine ( <i>Ipomea plebeian</i> ) *	1400	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 280 mL/ha <sup>B</sup> - Terrad <sup>®</sup> or <sup>®</sup> 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L to increase the speed of brownout and may also improve final control - Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control - Comet <sup>®</sup> 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of weeds pre-flowering.
Bent grasses ( <i>Agrostis</i> species)	2300	<b>TIMING</b> Apply to actively growing plants in late Spring when they have some seed-head development, but before Summer moisture stress. Remove stock to ensure there is full leaf growth. <b>FOLLOW-UP MANAGEMENT</b> Full disturbance with a tined implement should follow 10-21 days after spraying. Then follow with a Summer crop, and/or re-seeded pasture or crop the following Autumn.
Bifora ( <i>Bifora testiculata</i> )	2300	Nail 600EC Herbicide must always be added at 10-30 mL/ha.
Billygoat weed ( <i>Ageratum</i> species)	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 390 mL/ha <sup>B</sup>
Black bindweed ( <i>Fallopia convolvulus</i> )	1280	Alternatively to Nufarm weedmaster DST alone add either: - Kamba 750 at 105-160 mL/ha to improve control - Comet 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of 2 to 10 leaf weeds (up to 20 cm diameter) OR 375 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of 10 to 12 leaf weeds (up to 30 cm diameter).
Black spear grass ( <i>Heteropogon contortus</i> )	2300	
Blackberry nightshade ( <i>Solanum nigrum</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>

WEEDS	MINIMUM RATE (mL/ha) <sup>A</sup>	CRITICAL COMMENTS
Bladder ketmia ( <i>Hibiscus trionum</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 745 mL/ha<sup>B</sup></li> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant<sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control</li> <li>- Comet 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of 4 to 8 leaf weeds (up to 10 cm diameter).</li> </ul>
Blady grass ( <i>Imperata cylindrica</i> )	7000	
Blue top ( <i>Ageratum</i> species)	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 390 mL/ha <sup>B</sup>
Boggabri weed ( <i>Amaranthus mitchellii</i> )	770	
Bracken ( <i>Pteridium esculentum</i> )	7000	For Bracken add Pulse <sup>®</sup> at 200-500 mL/100L spray mix.
Brome grasses ( <i>Bromus</i> species)	960	Alternatively to Nufarm weedmaster DST alone add Terrad'or <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control.
Burr medic ( <i>Medicago polymorpha</i> )	960	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 390 mL/ha <sup>B</sup>
Button grass ( <i>Dactyloctenium radulans</i> )	770	
Californian burr ( <i>Xanthium orientale</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Calomba daisy ( <i>Oncosiphon suffruticosum</i> )	770	
Caltrop ( <i>Tribulus terrestris</i> ) *	770	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 280 mL/ha<sup>B</sup></li> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant<sup>^</sup> at 1 L/100L to increase the speed of brownout and may also improve final control</li> <li>- Kamba 750 at 105-160 mL/ha</li> <li>- Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control</li> <li>- Comet 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of weeds up to 15 cm diameter.</li> </ul>
Camel/Afghan melon ( <i>Citrullus lanatus</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 745 mL/ha<sup>B</sup></li> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 40 g/ha with Banjo Spray Adjuvant<sup>^</sup> at 1 L/100L to increase the speed of brownout and may also improve final control.</li> </ul>
Canary grass (Annual Phalaris) ( <i>Phalaris canariensis</i> )	960	
Capeweed ( <i>Arctotheca calendula</i> )	960	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant<sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Kamba 750 at 105-160 mL/ha for improved control</li> <li>- Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control</li> <li>- Nail 600EC Herbicide at 10-30 mL/ha to increase the speed of brownout and may also improve final control.</li> </ul>
Carpet grasses ( <i>Axonopus</i> species)	2300	
Charlock ( <i>Sinapsis arvensis</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 390 mL/ha <sup>B</sup>
Chickweed ( <i>Stellaria media</i> )	1530	Alternatively to Nufarm weedmaster DST alone add Nail 600EC Herbicide at 10-30 mL/ha for improved control.
Clammy Goosefoot ( <i>Chenopodium pumilio</i> )	770	Always add Terrad'or <sup>®</sup> 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L.
Climbing buckwheat ( <i>Fallopia convolvulus</i> )	1280	Alternatively to Nufarm weedmaster DST alone improved control occurs by adding either: <ul style="list-style-type: none"> <li>- Kamba 750 at 105-160 mL/ha</li> <li>- Bromicide<sup>®</sup> 200 at 1.5 L/ha .</li> </ul>
Clover species ( <i>Trifolium</i> species)	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 515 mL/ha<sup>B</sup></li> <li>- Kamba 750 at 270 mL/ha</li> <li>- Archer 750 Dual Salt Liquid Herbicide at 60-120 mL/ha.</li> </ul>
Cobblers peg ( <i>Bidens pilosa</i> )	1530	
Cocksfoot ( <i>Dactylis glomerata</i> )	2300	
Columbus grass ( <i>Sorghum almum</i> ) *, #	770	

WEEDS	MINIMUM RATE (mL/ha) <sup>A</sup>	CRITICAL COMMENTS
Common ice plant ( <i>Mesembryanthemum crystallinum</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Common sowthistle ( <i>Sonchus oleraceus</i> )	770	Previously grazed plants may be difficult to control without allowing full recovery. Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 280 mL/ha <sup>B</sup> - Comet 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of weeds at 2 to 5 leaf (up to 10 cm diameter).
Couch ( <i>Cynodon dactylon</i> ) #	2300	Best control of couch in WA and SA is obtained with Spring treatment. Most effective control of couch in eastern states is obtained with Summer and Autumn treatments. In cultivated situations use sequential treatments of 1.8-3.8 L/ha for control, otherwise use the maximum rate of 5.5 L/ha.
Cowvine ( <i>Ipomoea lonchophylla</i> )	1400	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup> - Terrad <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control - Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control - Comet 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of 2 to 10 leaf weeds (up to 10 cm diameter).
Crabgrass ( <i>Digitaria ciliaris</i> )	770	
Cudweeds ( <i>Gnaphalium</i> species)	770	
Dead nettle ( <i>Lamium amplexicaule</i> )	770	In Winter (cold) conditions, symptoms may be slow to develop.
Docks ( <i>Rumex</i> species) *	770	Established plants require minimum rate of Nufarm weedmaster DST of 1.92 L/ha. Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 390 mL/ha <sup>B</sup> - Kamba 750 at 270 mL/ha.
Doublegee ( <i>Emex australis</i> )	770	Alternatively to Nufarm weedmaster DST alone add either: - Terrad <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L to increase the speed of brownout and may also improve final control - Kamba 750 at 105-160 mL/ha for improved control - Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control - Nail 600EC Herbicide at 10-30 mL/ha to increase the speed of brownout and may also improve final control.
English couch ( <i>Elytrigia repens</i> )	4600	
Erodium ( <i>Erodium</i> species) *	1150	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup> - Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control.
False castor oil plant ( <i>Datura stramonium</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup> - Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control.
Fat hen ( <i>Chenopodium album</i> )	1400	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup> - Terrad <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control.
Fescues ( <i>Vulpia</i> species) #	1150	When treating dense infestations, nozzles designed to give MEDIUM to COARSE spray quality and a spray volume of 70 L/ha or more are recommended to improve spray coverage. Good coverage critical for control.
Flatweed ( <i>Hypochaeris radiata</i> )	1900	
Forage rape ( <i>Brassica campestris spp</i> )	770	Always add Terrad <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L (or other registered high quality methylated seed oil adjuvant).
Fumitories ( <i>Fumaria</i> species)	960	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 390 mL/ha <sup>B</sup> - Terrad <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L to increase the speed of brownout and may also improve final control.
Green Amaranth ( <i>Amaranthus viridis</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup>
Ground cherry ( <i>Physalis angulata</i> )	1530	
Guinea grass ( <i>Megathyrsus maximus</i> )	7000	
Hedge mustard ( <i>Sisymbrium officinale</i> )	770	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 280 mL/ha <sup>B</sup>

WEEDS	MINIMUM RATE (mL/ha) <sup>A</sup>	CRITICAL COMMENTS
Heliotrope ( <i>Heliotropium europaeum</i> )	770	Always add Terrad <sup>®</sup> or <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>†</sup> at 1 L/100L.
Hexham scent ( <i>Mellilotus indica</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Hoary cress ( <i>Lepidium draba</i> )	1150	Treat from late rosette to early flowering.
Horehound ( <i>Marrubium vulgare</i> ) *	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Indian hedge mustard ( <i>Sisymbrium orientale</i> )	770	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 280 mL/ha <sup>B</sup>
Ivyleaf speedwell ( <i>Veronica hederifolia</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Johnson grass ( <i>Sorghum halepense</i> ) #	4600	Apply to plants with a minimum of 30 cm new growth. Sequential treatments will be required for long term control.
Kangaroo grass ( <i>Themeda triandra</i> )	4600	
Kikuyu ( <i>Pennisetum clandestinum</i> )	4600	
Lesser canary grass ( <i>Phalaris minor</i> )	960	
Lesser swinecress ( <i>Lepidium didymum</i> )	2300	
Lincoln weed ( <i>Diplotaxis tenuifolia</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Liverseed grass ( <i>Urochloa panicoides</i> ) #	770	Under Summer (hot) conditions, dense infestations may require follow-up treatment for complete control. <b>DO NOT</b> tank mix with atrazine when spraying. Alternatively to Nufarm weedmaster DST alone add Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control.
Lovegrasses ( <i>Eragrostis</i> species) #	2300	
Marshmallow ( <i>Malva parviflora</i> ) *	1400	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup> - Terrad <sup>®</sup> or <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>†</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control - Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control - Nail 600EC Herbicide at 10-30 mL/ha to increase the speed of brownout and may also improve final control.
Medics ( <i>Medicago</i> species)	960	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 390 mL/ha <sup>B</sup> - Terrad <sup>®</sup> or <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>†</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control - Kamba 750 at 105-160 mL/ha - Archer 750 Dual Salt Liquid Herbicide at 60-120 mL/ha - Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control.
Mellilotus ( <i>Mellilotus indica</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Melons (Various genera)	1150	See also, Prickly paddy melon. Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup>
Mexican poppy ( <i>Argemone ochroleuca</i> )	770	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup>
Mintweed ( <i>Salvia reflexa</i> )	770	Alternatively to Nufarm weedmaster DST alone add Kamba 750 at 105-160 mL/ha for improved control.
Native millet ( <i>Panicum decompositum</i> ) #	770	
Native rosella ( <i>Hibiscus heterophyllus</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup>
Native vinya ( <i>Vigna lanceolata</i> )	770	Always add Terrad <sup>®</sup> or <sup>®</sup> 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant <sup>†</sup> at 1 L/100L to increase the speed of brownout and may also improve final control.
Navua sedge ( <i>Cyperus aromaticus</i> )	2300	
New Zealand spinach ( <i>Tetragonia tetragonoides</i> ) *	770	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 280 mL/ha <sup>B</sup>
Noogoora burr ( <i>Xanthium strumarium</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup> - Kamba 750 at 105-160 mL/ha - Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control.
Nutgrass ( <i>Cyperus rotundus</i> )	2300 followed by a second application of 2300	Apply two treatments of Nufarm weedmaster DST. Make first application to actively growing plants when at least 20% have reached the head stage (normally about Feb). After allowing maximum re-emergence to occur (normally in 6-8 weeks), it is essential to make a second application. NOTE Follow up treatments should be made as part of a Nutgrass control program.
Paragrass ( <i>Urochloa mutica</i> )	7000	Use on Dry Drains and Channels ONLY. <b>DO NOT</b> use in aquatic situations.

WEEDS	MINIMUM RATE (mL/ha) <sup>A</sup>	CRITICAL COMMENTS
Paradoxa grass ( <i>Phalaris paradoxa</i> )	960	
Paspalum ( <i>Paspalum dilatatum</i> )	4600	
Paterson's curse ( <i>Echium plantagineum</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 390 mL/ha<sup>B</sup></li> <li>- Kamba 750 at 105-160 mL/ha for improved control</li> <li>- Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control</li> <li>- Nail 60DEC Herbicide at 10-30 mL/ha to increase the speed of brownout and may also improve final control.</li> </ul>
Phalaris ( <i>Phalaris</i> species) #	1150	Nufarm weedmaster DST will provide knockdown, seasonal suppression and reduction in treated plant numbers. Alternatively to Nufarm weedmaster DST alone add Terrad'or <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>†</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control.
Pigweed, black/giant ( <i>Trianthema portulacastrum</i> )	770	
Pigweed, red ( <i>Portulaca oleracea</i> )	770	Plants up to 20 cm diameter will be controlled with an increased rate of at least 1200 mL/ha. Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Associate<sup>®</sup> at 7 g/ha for control over dense weed populations up to 6-leaf</li> <li>- Comet 400 Herbicide at 185 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of weeds up to 10 cm diameter OR 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of weeds up to 60 cm diameter.</li> </ul>
Plantains ( <i>Plantago</i> species)	1150	
Poa tussock ( <i>Poa labillardierei</i> ) (suppression)	2300	For reduction of ground cover allowing pasture renovation TIMING: Graze heavily, then remove at least 14 days before spraying to allow fresh regrowth. Apply to actively growing plants after the Autumn break but before heavy frosts (March – May). APPLICATION: Increasing to the higher rate may give more effective reductions. FOLLOW UP MANAGEMENT: Sowing may start from 14 days after spraying. It is essential that correct follow-up crop/pasture establishment and management occurs after each treatment. Spot treatment will limit re-infestation.
Polymeria, annual ( <i>Polymeria pusilla</i> )	1150	Alternatively to Nufarm weedmaster DST alone add Comet 400 Herbicide at 250 mL/ha for control of weeds 2 to 10 leaf (up to 20 cm diameter).
Potato weed ( <i>Galinsoga parviflora</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup>
Prairie grass ( <i>Bromus catharticus</i> )	4600	
Prickly lettuce ( <i>Lactuca serriola</i> )	1400	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 390 mL/ha<sup>B</sup></li> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant<sup>†</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Kamba 750 at 105-160 mL/ha</li> <li>- Comet 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of 2 to 5 leaf weeds (up to 10 cm diameter).</li> </ul>
Prickly paddy melon ( <i>Cucumis myriocarpus</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 745 mL/ha<sup>B</sup></li> <li>- a triclopyr 600 g/L product at 80 mL/ha plus Nufarm weedmaster DST at 770 mL/ha. <b>DO NOT</b> add crop oil.</li> </ul>
Quack grass ( <i>Elytrigia repens</i> )	4600	
Queensland Bluegrass ( <i>Dichanthium sericeum</i> )	4600	
Raspweed ( <i>Haloragis aspera</i> ) *	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 280 mL/ha <sup>B</sup>
Red-leg grass ( <i>Bothriochloa macra</i> )	4600	
Redshank ( <i>Pericaria maculosa</i> )	770	Kamba 750 must always be added at 105-160 mL/ha.
Red Natal grass ( <i>Melinis repens</i> )	2300	
Rhodes grass ( <i>Chloris gayana</i> )	4600	
Rhynchosia ( <i>Rhynchosia minima</i> )	1150	Comet 400 Herbicide must always be added at 250 mL/ha for control of weeds at seedling to early flowering.
Rope twitch ( <i>Elytrigia repens</i> )	4600	
Ryegrasses ( <i>Lolium</i> species)	1150	



WEEDS	MINIMUM RATE (mL/ha) <sup>A</sup>	CRITICAL COMMENTS
Saffron thistle ( <i>Carthamus lanatus</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 390 mL/ha<sup>B</sup></li> <li>- Terrad<sup>or</sup>® 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant<sup>A</sup> at 1 L/100L to increase the speed of brownout and may also improve final control.</li> </ul>
Scotch thistle ( <i>Onopordum acanthium</i> )	1150	
Serrated Tussock ( <i>Wassella trichotoma</i> )	3100	Apply to actively growing and stress-free plants. Best results May to October. <b>Application:</b> Boom spray volume of 70 L/ha or more is recommended to improve plant coverage. Also see Aerial Equipment. <b>Surfactants:</b> Addition of 200 mL of Wetter TX™ to 100 L of spraying solution may improve control of Serrated Tussock. <b>Site Preparation:</b> <i>Burning</i> of Serrated Tussock 10-12 months before spraying or <i>slashing/heavy grazing</i> (cell grazing) 2 weeks before spraying is essential for good results ( <b>Note:</b> Serrated Tussock is almost indigestible and prolonged exposure can lead to starvation and death of stock). <b>Rates:</b> Use the 3.1 L/ha rate on Serrated Tussock regrowth after burning (no residual dead foliage). Use the 7.0 L/ha rate on Serrated Tussock that has been slashed or grazed (may contain some residual dead foliage).
Sesbania pea ( <i>Sesbania cannabina</i> )	1150	Comet 400 Herbicide must always be added at 250 mL/ha for control of weeds at 2 to 6 leaf (up to 10 cm tall).
Shepherd's purse ( <i>Capsella bursa-pastoris</i> )	1400	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 515 mL/ha<sup>B</sup></li> <li>- Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control.</li> </ul>
Silver grasses ( <i>Vulpia</i> species) #	1150	When treating dense infestations, nozzles designed to give MEDIUM to COARSE spray quality and a spray volume of 70 L/ha or more are recommended to improve spray coverage. Good coverage critical for control. Alternatively to Nufarm weedmaster DST alone add Terrad <sup>or</sup> ® 700WG Herbicide at 30-40 g/ha with Banjo Spray Adjuvant <sup>A</sup> at 1 L/100L to increase the speed of brownout and may also improve final control.
Silver-leaf nightshade ( <i>Solanum elaeagnifolium</i> )	7000	
Skeleton weed ( <i>Chondrilla juncea</i> ) *	1150	Nufarm weedmaster DST will provide knockdown, seasonal <b>suppression</b> and reduction in treated plant numbers only. Plants need to be fully emerged rosettes (NSW only). Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Soldier thistle ( <i>Picnomon acarna</i> )	1400	Archer 750 Dual Salt Liquid Herbicide must always be added at 120 mL/ha for control of weeds up to 10cm diameter at the 4 to 8 leaf stage.
Sorrel ( <i>Rumex acetosella</i> ) *	1150	Nufarm weedmaster DST will provide knockdown, seasonal suppression and reduction in treated plant numbers. Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 515 mL/ha<sup>B</sup></li> <li>- Terrad<sup>or</sup>® 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant<sup>A</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Kamba 750 at 270 mL/ha.</li> </ul>
Soursob ( <i>Oxalis pes-caprae</i> )	1150	Nufarm weedmaster DST will provide knockdown, seasonal suppression and reduction in treated plant numbers. Treat repeatedly until tuber exhaustion.
Sow thistle ( <i>Sonchus oleraceus</i> )	770	Previously grazed plants may be difficult to control without allowing full recovery. Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 280 mL/ha<sup>B</sup></li> <li>- Terrad<sup>or</sup>® 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant<sup>A</sup> at 1 L/100L to increase the speed of brownout and may also improve final control</li> <li>- Kamba 750 at 105-160 mL/ha</li> <li>- Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control.</li> </ul>
Spear thistle ( <i>Cirsium vulgare</i> )	1150	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 390 mL/ha <sup>B</sup>
Spineless caltrop ( <i>Tribulus micrococcus</i> )	1150	Comet 400 Herbicide must always be added at 250 mL/ha for control of weeds up to 15 cm diameter.
Spiny burgrass ( <i>Cenchrus longispinus</i> )	1530	
Spurges ( <i>Eupatorium</i> species)	770	
Stinkgrass ( <i>Eragrostis cilianensis</i> ) #	770	
Stinking goosefoot ( <i>Chenopodium vulvaria</i> )	770	

WEEDS	MINIMUM RATE (mL/ha) <sup>a</sup>	CRITICAL COMMENTS
Storksbills ( <i>Erodium</i> spp) *	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 515 mL/ha<sup>a</sup></li> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant<sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Nail<sup>®</sup> 600EC Herbicide at 10-30 mL/ha for improved control of weeds with a maximum of 4 leaves.</li> </ul>
Subterranean clover ( <i>Trifolium subterraneum</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 515 mL/ha<sup>a</sup></li> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant<sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Kamba 750 at 270 mL/ha</li> <li>- Archer 750 Dual Salt Liquid Herbicide at 60-120 mL/ha</li> <li>- Nail 600EC Herbicide at 10-30 mL/ha for improved control.</li> </ul>
Summer grass ( <i>Digitaria ciliaris</i> )	770	
Sweet summer grass ( <i>Brachiaria eruciformis</i> )	770	
Thornapples ( <i>Datura</i> species)	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 515 mL/ha<sup>a</sup></li> <li>- Kamba 750 at 105-160 mL/ha</li> <li>- Comet 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of 2 to 8 leaf weeds (up to 15 cm diameter).</li> </ul>
Tree hogweed ( <i>Polygonum patulum</i> )	770	Kamba 750 must always be added at 105-160 mL/ha.
Turnip weed ( <i>Rapistrum rugosum</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 280 mL/ha<sup>a</sup></li> <li>- Kamba 750 at 105-160 mL/ha.</li> </ul>
Tussock grass ( <i>Poa labillardierei</i> )	2300	See Poa tussock.
Variegated thistle ( <i>Silybum marianum</i> )	960	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 390 mL/ha<sup>a</sup></li> <li>- Kamba 750 at 105-160 mL/ha.</li> </ul>
Volunteer beans (Various species)	2300	Always add either: <ul style="list-style-type: none"> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 15-40 g/ha (refer to Terrad'or<sup>®</sup> label for rates on each volunteer bean) with Banjo Spray Adjuvant<sup>^</sup> at 1 L/100L to increase the speed of brownout and may also improve final control</li> <li>- Archer 750 Dual Salt Liquid Herbicide at 60-120 mL/ha.</li> </ul>
Volunteer canola ( <i>Brassica napus</i> ) including conventional, Roundup Ready <sup>®</sup> varieties and canola hybrids with the Optimum GLY <sup>®</sup> herbicide tolerance trait (herbicide tolerant varieties only when tank mixed with Amicide Advance 700 or Terrad'or <sup>®</sup> 700WG Herbicide)	1400	Always use 900 mL/ha of Amicide Advance 700 up to the 4 leaf weed stage. Use 1500 mL/ha of Amicide Advance 700 up to the 6 leaf weed stage <sup>®</sup> . For adequate coverage use a minimum application water volume of 70 L/ha. It is essential to manage volunteer herbicide tolerant canola varieties in both crop and non-crop situations. These plants are best managed using an integrated weed management program that may include the use of alternative mode of action herbicides registered for use in the crop and/or cultivation. Alternatively to Amicide Advance 700 add Terrad'or <sup>®</sup> 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control
Volunteer cereals (Various species)	770	Alternatively to Nufarm weedmaster DST alone add Terrad'or <sup>®</sup> 700WG Herbicide at 15-40 g/ha (refer to Terrad'or <sup>®</sup> label for rates on each volunteer cereal) with Banjo Spray Adjuvant <sup>^</sup> at 1 L/100L to increase the speed of brownout and may also improve final control.
Volunteer cotton ( <i>Gossypium</i> spp.), seedlings, including Roundup Ready <sup>®</sup> varieties	2300	Always add either: <ul style="list-style-type: none"> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 40 g/ha with Banjo Spray Adjuvant<sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Nail 600EC Herbicide at 30-40 mL/ha (for Roundup Ready<sup>®</sup> varieties) or 20-30 mL/ha (for conventional varieties) and apply with Supercharge<sup>®</sup> Elite or Banjo<sup>®</sup> at 0.5 L/100L</li> <li>- Bromicide 200 Herbicide at 1.5 L/ha.</li> </ul>
Volunteer lentils ( <i>Lens culinaris</i> )	1400	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Terrad'or<sup>®</sup> 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant<sup>^</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Archer 750 Dual Salt Liquid Herbicide at 60-120 mL/ha</li> <li>- Kamba 750 at 105-160 mL/ha.</li> </ul>

WEEDS	MINIMUM RATE (mL/ha) <sup>A</sup>	CRITICAL COMMENTS
Volunteer lupins ( <i>Lupinus</i> species)	960	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 390 mL/ha<sup>B</sup></li> <li>- Terrad<sup>or</sup>® 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant<sup>C</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Archer 750 Dual Salt Liquid Herbicide at 60-120 mL/ha</li> <li>- Kamba 750 at 105-160 mL/ha.</li> </ul>
Volunteer peas (Various species)	1400	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 390 mL/ha<sup>B</sup></li> <li>- Terrad<sup>or</sup>® 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant<sup>C</sup> at 1 L/100L to increase the speed of brownout and may also improve final control</li> <li>- Archer 750 Dual Salt Liquid Herbicide at 60-120 mL/ha</li> <li>- Kamba 750 at 105-160 mL/ha.</li> </ul>
Volunteer safflower ( <i>Carthamus tinctorius</i> )	1400	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 515 mL/ha<sup>B</sup></li> <li>- Archer 750 Dual Salt Liquid Herbicide at 120 mL/ha for control of weeds up to 6 leaf.</li> </ul>
Volunteer sorghum ( <i>Sorghum bicolor</i> )	770	
Volunteer sunflower ( <i>Helianthus annuus</i> )	770	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 390 mL/ha<sup>B</sup></li> <li>- Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100L to increase the speed of brownout and may also improve final control.</li> </ul>
Volunteer vetch ( <i>Vicia sativa</i> )	1400	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 515 mL/ha<sup>B</sup></li> <li>- Terrad<sup>or</sup>® 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant<sup>C</sup> at 1 L/100L to increase the speed of brownout and may also improve final control</li> <li>- Archer 750 Dual Salt Liquid Herbicide at 60-120 mL/ha.</li> </ul>
Ward's weed ( <i>Carrichtera annua</i> )	1400	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 515 mL/ha <sup>B</sup>
Water couch ( <i>Paspalum distichum</i> )	7000	Use on Dry Drains and Channels ONLY. <b>DO NOT</b> use in aquatic situations.
Wild gooseberry ( <i>Physalis angulata</i> )	1150	Alternatively to Nufarm weedmaster DST alone add Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup>
Wild lettuce ( <i>Lactuca saligna</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 390 mL/ha<sup>B</sup></li> <li>- Kamba 750 at 105-160 mL/ha.</li> </ul>
Wild mustard ( <i>Sisymbrium officinale</i> )	1150	Alternatively to Nufarm weedmaster DST alone add Kamba 750 at 105-160 mL/ha.
Wild oats ( <i>Avena</i> species)	770	Alternatively to Nufarm weedmaster DST alone add Terrad <sup>or</sup> ® 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant <sup>C</sup> at 1 L/100L to increase the speed of brownout and may also improve final control.
Wild radish ( <i>Raphanus raphanistrum</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 280 mL/ha<sup>B</sup></li> <li>- Terrad<sup>or</sup>® 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant<sup>C</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> <li>- Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100 L to increase the speed of brownout and may also improve final control</li> <li>- Nail 600EC Herbicide at 10-30 mL/ha to increase the speed of brownout and may also improve final control.</li> </ul>
Wild turnip ( <i>Brassica tournefortii</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 280 mL/ha<sup>B</sup></li> <li>- Terrad<sup>or</sup>® 700WG Herbicide at 15-40 g/ha with Banjo Spray Adjuvant<sup>C</sup> at 1 L/100L and 0.77-2.3 L/ha of Nufarm weedmaster DST to increase the speed of brownout and may also improve final control</li> </ul>
Winter grass ( <i>Poa annua</i> )	960	Alternatively to Nufarm weedmaster DST alone add Terrad <sup>or</sup> ® 700WG Herbicide at 20-40 g/ha with Banjo Spray Adjuvant <sup>C</sup> at 1 L/100L to increase the speed of brownout and may also improve final control
Wireweed ( <i>Polygonum aviculare</i> )	1150	Alternatively to Nufarm weedmaster DST alone add either: <ul style="list-style-type: none"> <li>- Amicide Advance 700 at a minimum of 515 mL/ha<sup>B</sup></li> <li>- Kamba 750 at 105-160 mL/ha</li> <li>- Terrain 500 WG Herbicide at 30 g/ha with Banjo Spray Adjuvant at 0.5-1 L/100 L to increase the speed of brownout and may also improve final control</li> <li>- Comet 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of weeds at 2 to 3 leaf (up to 10 cm tall).</li> </ul>
Wire grasses ( <i>Aristida</i> species)	2300	

WEEDS	MINIMUM RATE (mL/ha) <sup>A</sup>	CRITICAL COMMENTS
Yellow burr weed ( <i>Amsinckia calycina</i> )	960	
Yellow vine ( <i>Tribulus micrococcus</i> ) <sup>*</sup>	770	Alternatively to Nufarm weedmaster DST alone add either: - Amicide Advance 700 at a minimum of 745 mL/ha <sup>B</sup> - Kamba 750 at 105-160 mL/ha. - Comet 400 Herbicide at 250 mL/ha plus 1150 mL/ha Nufarm weedmaster DST for control of weeds up to 15 cm diameter.
Yorkshire fog ( <i>Holcus lanatus</i> ) #	1150	

# Addition of Wetter TX 200 mL/100L spray solution may improve control      \* Plants of this species are best controlled as seedlings

<sup>^</sup> Or other registered high quality methylated seed oil adjuvant

#### GENERAL COMMENTS FOR SECTION 2

##### <sup>A</sup> Application rate selection:

Use Nufarm weedmaster DST at 0.6-5.5 L/ha as directed. Minimum application rates for each weed are listed in the table above (in mL/ha); recommended use rates (except where a tank mixture is used where the maximum rate is 2.3 L/ha, or when applying by aircraft when the maximum rate is 3.1 L/ha) are:

- 0.77-2.3 L/ha for annual weed species
- 2.3-64.6 L/ha for perennial species
- 7.0 L/ha for hard to control perennial species

NOTE: The maximum approved rate for any weed is 7.0 L/ha and this should never be exceeded.

Use lower rates on young weeds and where cultivation is to follow within 21 days. Increase to higher rates when treating under cold/overcast conditions, or when treating late in the season, or where grasses reach full tillering, or where broadleaf weeds reach stem elongation/budding, or when commencing a fallow. Addition of Liase at 2 L/100 L, may improve control when treating under adverse environmental conditions.

##### <sup>B</sup> Perennial weeds:

Control of established perennials is best obtained when plants are at the seedhead stage. In general, best control of Winter growing perennials is obtained with application during Winter-Spring. Best control of Summer growing perennials is obtained with application late Summer and Autumn.

##### Tank mixtures:

See **TANK MIXTURES** for directions. Read and follow all label directions, restraints, plant-back periods, withholding periods, regional use restrictions and safety directions for the tank mix products. Nufarm weedmaster DST always requires the addition of a tank-mix surfactant such as Activator or Collide 700 (refer to the Surfactant Addition section of the General Instructions), except where specifically noted in the Directions for Use. Use Collide 700 with the mixture product if insecticides will be included in the tank mixture or if faster brown out of weeds is required. Additional surfactant requirements are listed for specific tank-mix partners in the table above. For residual control of annual weeds Nufarm weedmaster DST may be tank-mixed with certain residual herbicides listed in **TANK MIXTURES**.

**DO NOT** use Nufarm weedmaster DST at rates higher than 2.3 L/ha in a tank mix for uses in Section 1 unless otherwise stated in the Section 2 weed table above.

##### Aerial application:

Use Nufarm weedmaster DST up to a maximum of 3.1 L/ha. For further instructions on aerial application, and application under hot conditions, see **AERIAL EQUIPMENT**.

##### <sup>B</sup> Amicide Advance 700 rate selection:

Use Amicide Advance 700 at 0.28-1.2 L/ha as directed in a tank mixture with Nufarm weedmaster DST. Minimum application rates for each weed are listed above, however all weeds listed can be treated up to the maximum application rate of 2.3 L/ha of Nufarm weedmaster DST and 1.2 L/ha of Amicide Advance 700. Use lower rates for seedling broadleaf weeds and increase to higher rates for broadleaf weeds more than 10 cm diameter/high. For rate ranges specific to individual species, see the table above. At the time of application, all weeds must be actively growing and not under stress from low moisture, frost, cold, disease or water-logging. If grazing has occurred allow regrowth to 6-8 cm before spraying and use higher rate. For adequate coverage use a minimum application water volume of 70 L/ha.

Section 3. FLEABANE & SOWTHISTLE CONTROL FOR FALLOW AND PRE-SOWING – SEQUENTIAL APPLICATION

SITUATION & CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
<p>Prior to sowing any crop or pasture or for a fallow (refer to partner product labels for planting intervals)</p>	<p>Flaxleaf fleabane (<i>Coryza bonariensis</i>) Sowthistle (<i>Conchus oleraceus</i>)</p>	<p>First spray application:  1.4-2.3 L/ha + 0.65-1.1 L/ha Amicide Advance 700</p>	<p>Apply to cotyledon to 12 leaf rosette prior to stem elongation. For Amicide Advance 700 use the low rate in Autumn/Winter and use the highest rate for Spring/Summer applications. For adequate coverage use a minimum application water volume of 70 L/ha. A sequential application of Shirquat or Terrad'or® 700WG Herbicide (refer below) is also recommended for situations where incomplete control is achieved with the first application, or where there are spray misses/shadowing, failures due to resistance, or under periods of temperature and/or moisture stress. In these situations, the sequential application is to be applied 7-14 days after the first application. <b>DO NOT</b> apply prior to transplanting tomato seedlings. Refer to Amicide Advance 700 label for planting intervals.</p>
		<p>Sequential spray application:  1.6-2.0 L/ha Shirquat*  OR  40 g/ha Terrad'or® 700WG Herbicide + 1% v/v Banjo Spray Adjuvant*</p>	<p><u>Second application with Shirquat:</u> Apply at stem elongation to flowering plants. Apply the sequential application 7-14 days after the first application. For Shirquat use the low rate in Autumn/Winter and use the highest rate for Spring/Summer applications. For adequate coverage use a minimum application water volume of 70 L/ha. The sequential application of Shirquat is recommended for situations where incomplete control is achieved with the first application, or where there are spray misses/shadowing, failures due to resistance or under periods of temperature and/or moisture stress. In these situations, the sequential application is to be applied 7-14 days after the first application.</p> <p><u>Second application with Terrad'or® 700WG Herbicide:</u> Apply the sequential application (follow up spray) 7-14 days after the first application. For adequate coverage use a minimum application water volume of 80 L/ha. Observe the plant-back periods listed in the GENERAL INSTRUCTIONS. Sow following crops with a seeder that will move treated soil away from crop row (e.g. knife point with press wheels). Use of seeding machinery, or planting under conditions where product or treated soil remains or moves back into the crop row may result in crop damage. This may be caused by factors such as inappropriate seeding machinery, narrow row spacings, fast travel speeds, soil type, wind, heavy rainfall or irrigation after planting, etc. This is particularly important for susceptible crops such as canola. <b>DO NOT</b> apply post-sowing pre-emergent. Addition of Banjo spray adjuvant* at 1.0 % v/v must always be used with Terrad'or® 700 WG. Rainfast after 1 hour of daylight after application to seedling annual weeds when used with Banjo (for other mixtures, refer to tank mix partner label for rainfastness information). To broaden the weed spectrum Terrad'or® 700WG Herbicide may be tank mixed with the recommended rate of another knockdown herbicide (Shirquat®, Revolver®, Biffo®, Amitrole T). * or other registered high quality methylated seed oil adjuvant</p>

#### Section 4. CANOLA - HYBRIDS WITH THE OPTIMUM GLY® HERBICIDE TOLERANCE TRAIT

##### CROP SAFETY

Over-the-top (OTT) applications may be made in canola hybrids with the Optimum GLY® Herbicide Tolerance Trait from cotyledon (BBCH 10) to the early bloom (BBCH 61) stage. Sequential applications must be at least 14 days apart.

The canola crop must not have advanced beyond the latest recommended growth stage (i.e. 10% of flowers on main raceme open, main raceme elongating).

No additional surfactant is required for use in canola hybrids with the Optimum GLY® Herbicide Tolerance Trait.

NOTE: The total glyphosate active constituent applied as OTT applications up to BBCH61 in any one crop must not exceed 3.24 kg/ha.

Before application, users should consult the Optimum GLY® Technical Manual which have been designed to minimise the development of glyphosate resistance in weed populations.

WEEDS CONTROLLED	GROWTH STAGE OF CROP	RATE	WHP	CRITICAL COMMENTS	
African turnip weed, Amsinckia, Annual ryegrass, Australian bluebell, Barley grass, Brome grass, Canary grass, Capeweed, Chickweed, Climbing buckwheat (less than 12 leaves), Cudweed, Deadnettle, Dock (seedling), Doublegee, Erodium, Flatweed, Fumitory, Hoary cress (late rosette to early flowering), Indian hedge mustard, Lesser swinecress, Medic (annual), Mexican poppy, Milk thistle, Mintweed,	New Zealand spinach, Paradoxa grass, Paterson's curse, Prickly lettuce, Safron thistle, Scotch thistle, Silver grass, Skeleton weed (fully emerged rosettes), Spear thistle, Spiny Emex, Sub clover, Sow thistle, Volunteer cereals, Volunteer chickpeas, Volunteer field peas, Volunteer lentils, Volunteer Lupins, Variegated thistle, Wild mustard, Wild oats, Wild lettuce, Wild radish, Wild turnip, Winter grass, Wireweed	Cotyledon (BBCH 10 - Cotyledons completely unfolded) to early bloom (BBCH 61 - 10% of flowers on main raceme open, main raceme elongating)	1.3-2.3 L/ha	<p><b>Harvest:</b> Not required when used as directed.</p> <p><b>Grazing / Cutting for stockfeed:</b> 7 days</p>	<p><b>DO NOT</b> apply after BBCH 61 (10% of flowers on main raceme open, main raceme elongating) crop stage.</p> <p><b>DO NOT</b> apply more than 3 applications in any one crop. Each application must be a minimum of 1.3 L/ha and a maximum of 2.3 L/ha.</p> <p>Applications may be made to canola hybrids with the Optimum GLY® herbicide tolerance trait from cotyledon to the 10% flowering stage. The addition of a surfactant is not required for use in canola hybrids with the Optimum GLY® herbicide tolerance trait.</p> <p>Apply treatments to weeds which have at least one true leaf (broadleaf weeds) or two leaves (grasses) to provide an adequate surface area for herbicide uptake. Repeat applications may be required if a second flush of weeds germinate. For sequential applications, applications must be at least 14 days apart.</p> <p>Annual weeds are only to be sprayed when they are actively growing. Use the lower rate on grass weeds up to early tillering and broadleaf weeds with up to 2 true leaves. Increase to the higher rate on larger weeds and in situations of high density weed populations or tough environmental conditions.</p> <p>Perennial weeds should always be treated at the higher rate. Nufarm weedmaster DST will provide knockdown, seasonal suppression and reduction in treated plant numbers.</p> <p>The effects of this product may not be apparent for 3 to 7 days (annual weeds) or 2 to 3 weeks (perennial weeds) or longer under cool, cloudy conditions. This product will control emerged weeds only and provides no residual control.</p>
Weeds as above plus: Soldier thistle (up to 10 cm diameter, 4 to 8 leaf)*, Volunteer faba beans, Volunteer safflower (up to 6 leaf)*, Volunteer vetch	2 to 8 leaf (BBCH12 to BBCH18). One or two applications; however total rate of Archer 750 Dual Salt Liquid Herbicide should not exceed 120 mL/ha in the one season.	1.3 L/ha + 60-120 mL/ha Archer 750 Dual Salt Liquid Herbicide		<p>Use the higher rate of Archer 750 Dual Salt Liquid Herbicide in situations of high weed population, large weed size, and/or conditions of environmental stress (dry, frost etc).</p> <p>*The higher rate of Archer 750 Dual Salt Liquid Herbicide is required for control of soldier thistle and volunteer safflower.</p> <p>Varying levels of control can be experienced between different varieties of these species. Total application of Archer 750 Dual Salt Liquid Herbicide should not exceed 120 mL/ha in the one season. Application of this tank mix at the first spray timing is recommended.</p>	



**WARNING:** THE APPLICATIONS RECOMMENDED ABOVE ARE FOR USE ONLY WITH DESIGNATED CANOLA HYBRIDS WITH THE OPTIMUM GLY® HERBICIDE TOLERANCE TRAIT. APPLICATION OF THIS PRODUCT TO CANOLA THAT DOES NOT CONTAIN THE OPTIMUM GLY® HERBICIDE TOLERANCE TRAIT MAY RESULT IN SEVERE CROP INJURY AND YIELD LOSS OR DEATH OF THE CANOLA. EXTREME CARE MUST BE TAKEN TO AVOID CONTACT WITH CROPS OR DESIRABLE PLANTS WITHOUT THE OPTIMUM GLY® HERBICIDE TOLERANCE TRAIT, OR WITH NATIVE VEGETATION, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

**Section 5. CANOLA –TRUFLEX\* WITH ROUNDUP READY\* TECHNOLOGY**

**CROP SAFETY FOR OTT APPLICATIONS**

Over-the-top (OTT) applications may be made in canola varieties of TruFlex\* with Roundup Ready\* Technology (TruFlex\* Canola) from crop emergence to the first flowering stage. Sequential applications must be at least 14 days apart and the canola crop must have incremental growth of two leaves between applications. The canola crop must not have advanced beyond the latest recommended growth stage (i.e. first flowering).

No additional surfactant is required for use in TruFlex\* Canola.

NOTE: The total glyphosate active constituent applied as OTT applications up to first flowering in any one crop must not exceed 1.8 kg/ha (excluding pre-harvest applications).

**PRE-HARVEST**

Following OTT applications, Nufarm weedmaster DST can also be used pre-harvest, applied at early senescence. For details, please refer to Section 8.

<b>SITUATION - TRUFLEX* CANOLA</b>				
<b>Before use in this situation is carried out users should consult the TruFlex* Canola Resistance Management Plan (RMP) which has been developed to minimise the evolution of herbicide resistance in weed populations.</b>				
<b>WEEDS CONTROLLED</b>	<b>GROWTH STAGE OF CROP</b>	<b>GROWTH STAGE OF WEED</b>	<b>RATE</b>	<b>CRITICAL COMMENTS</b>
Annual ryegrass, Barley grass, Brome grass, Canary grass, Capeweed, Paterson's curse, Saffron thistle, Scotch thistle, Silver grass, Spear thistle, Variegated thistle, Volunteer cereals, Wild mustard, Wild oats, Wild radish, Wild turnip, Winter grass	Crop emergence to first flowering (at least 50% of plants have at least one flower).	For grass weeds and volunteer cereals: 1 leaf to mid-tillering  For volunteer plants and/ or broadleaf weeds: 1 true leaf to 8 leaves	1.3-1.9 L/ha	Up to 2 applications of up to 1.9 L/ha may be made in any one crop. Or, up to 3 applications of 1.3 L/ha may be made in any one crop. Repeat applications may be required if a second flush of weeds germinates but <b>DO NOT</b> apply after the first flowering stage of the crop (except when used pre-harvest – Refer to Section 8). For sequential applications, applications must be at least 14 days apart and the canola crop must have incremental growth of two leaves between applications. The canola crop must have not advanced beyond the latest recommended growth stage (i.e. first flowering). Ensure broadleaf weeds have at least one true leaf, and grasses two leaves before application. <b>DO NOT</b> apply after first flowering (except when used pre-harvest – Refer to Section 8).
Weeds as above plus, Annual medic, Sub clover, Volunteer chickpeas, Volunteer field peas, Volunteer lentils, Volunteer lupins	Crop emergence to first flowering (at least 50% of plants have at least one flower).  Two applications required.		1.3-1.9 L/ha	Two applications of Nufarm weedmaster DST provide higher levels of control than a single application. <b>DO NOT</b> apply after first flowering (except when used pre-harvest – Refer to Section 8).
Weeds as above plus, Volunteer faba beans, Volunteer vetch	Crop emergence to first flowering (at least 50% of plants have at least one flower).  One or two applications; however total rate of Archer 750 Dual Salt Liquid Herbicide should not exceed 120 mL/ha in the one season.		1.3-1.9 L/ha + 60-120 mL/ha Archer 750 Dual Salt Liquid Herbicide	Use the higher rate of Archer 750 Dual Salt Liquid Herbicide in situations of high weed population, large weed size, and/or conditions of environmental stress (dry, frost etc). Varying levels of control can be experienced between different varieties of these species. Total application of Archer 750 Dual Salt Liquid Herbicide should not exceed 120 mL/ha in the one season. Application of this tank mix at the first spray timing is recommended. <b>DO NOT</b> apply after first flowering (except when used pre-harvest – Refer to Section 8).

**WARNING: THE APPLICATIONS RECOMMENDED ABOVE ARE FOR USE ONLY WITH IMPROVED CANOLA VARIETIES THAT ARE DESIGNATED AS CANOLA WITH THE TRUFLEX\* TECHNOLOGY. SEVERE INJURY OR DEATH OF CANOLA WILL RESULT IF ANY CANOLA VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE TRUFLEX\* TECHNOLOGY ARE SPRAYED WITH THIS PRODUCT. EXTREME CARE MUST BE TAKEN TO AVOID CONTACT WITH CROPS OR DESIRABLE PLANTS WITHOUT THE TRUFLEX\* TECHNOLOGY, OR WITH NATIVE VEGETATION, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.**

**Section 6. CANOLA - ROUNDUP READY\***

**CROP SAFETY**

Over-the-top (OTT) applications may be made in Roundup Ready\* canola from crop emergence to the 6-leaf stage (prior to bud formation).

Sequential applications must be at least 14 days apart and canola must have incremental growth of at least 2 new leaves between applications.

**Some short-term, visual yellowing may occur when Nufarm weedmaster DST is applied. This effect is temporary and will not influence crop growth or yield.**

No additional surfactant is required for use in Roundup Ready\* canola varieties.

Nufarm weedmaster DST should be applied alone or with Archer 750 Dual Salt Liquid Herbicide, Nufarm Collide 700, Nufarm Liase, Astound Duo and Nufarm Dimethoate. Other tank mixes are not recommended for over-the-top applications of Nufarm weedmaster DST due to the potential for reduced weed control or crop injury. Nufarm Liase may increase the performance of this product on annual and perennial weeds, particularly under hard water conditions (high levels of calcium, magnesium or bicarbonate ions) or drought conditions.

**NOTE:** The total glyphosate active constituent applied as OTT applications up to the 6-leaf stage in any one crop must not exceed 1.2 kg/ha (excluding pre-harvest applications)

**PRE-HARVEST AID**

Nufarm weedmaster DST can be used pre-harvest, applied at early senescence. For details, please refer to Section 8.

**CAUTION:** The applications recommended here are for use only with improved canola varieties that are designated as glyphosate tolerant canola. Severe injury or death of canola will result if any canola varieties not properly designated as having glyphosate tolerance are sprayed with this product. Extreme care must be taken to avoid contact with crops or desirable plants without glyphosate tolerance, or with native vegetation, since severe injury or destruction will result.

<b>SITUATION - ROUNDUP READY* CANOLA</b> Before use in this situation is carried out users should consult the Resistance Management Plan which has been developed to minimise the evolution of herbicide resistance in weed populations.				
<b>WEEDS CONTROLLED</b>	<b>GROWTH STAGE OF CROP</b>	<b>GROWTH STAGE OF WEED</b>	<b>RATE</b>	<b>CRITICAL COMMENTS</b>
Annual ryegrass, Barley grass, Brome grass, Canary grass, Capeweed, Paterson's curse, Saffron thistle, Scotch thistle, Silver grass, Spear thistle, Variegated thistle, Volunteer cereals, Wild mustard, Wild oats, Wild radish, Wild turnip, Winter grass.	Crop emergence to 6 leaf (prior to bud formation).	For grass weeds and volunteer cereals: 1 leaf to mid-tillering. For volunteer plants and/or broadleaf weeds: 1 true leaf to 8 leaves.	1.3 L/ha	<b>Up to 2 applications only may be made in any one crop. Each application must be 1.3 L/ha.</b> Repeat applications may be required if a second flush of weeds germinates but <b>DO NOT</b> apply after the 6-leaf stage of the crop. For sequential applications, applications must be at least 14 days apart and the canola crop must have incremental growth of two leaves between applications. The canola crop must have not advanced beyond the latest recommended growth stage (i.e. 6 leaf). Ensure broadleaf weeds have at least one true leaf, and grasses two leaves before application.
Weeds as above plus: Annual Medic, Sub. clover, Volunteer Chickpeas, Volunteer Field peas, Volunteer Lentils, Volunteer Lupins	Crop emergence to 6 leaf (prior to bud formation). Two applications required		1.3 L/ha	Two applications of Nufarm weedmaster DST provide higher levels of control than a single application.
Weeds as above plus: Annual Medic, Sub. clover, Volunteer Faba beans, Volunteer Chickpeas, Volunteer Field peas, Volunteer Lentils, Volunteer Lupins, Volunteer Vetch	2 to 6 leaf (prior to bud formation). One or two applications; however total rate of Archer 750 Dual Salt Liquid Herbicide should not exceed 120 mL/ha in the one season.		1.3 L/ha + 60-120 mL/ha Archer 750 Dual Salt Liquid Herbicide	Use the higher rate of Archer 750 Dual Salt Liquid Herbicide in situations of high weed population, large weed size, and/or conditions of environmental stress (dry, frost etc). Varying levels of control can be experienced between different varieties of these species. Total application of Archer 750 Dual Salt Liquid Herbicide should not exceed 120 mL/ha in the one season. Application of this tank mix at the first spray timing is recommended.

**WARNING:** THE APPLICATIONS RECOMMENDED ABOVE ARE FOR USE ONLY WITH IMPROVED CANOLA VARIETIES THAT ARE DESIGNATED AS CANOLA WITH THE ROUNDUP READY\* TECHNOLOGY. SEVERE INJURY OR DEATH OF CANOLA WILL RESULT IF ANY CANOLA VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY\* TECHNOLOGY ARE SPRAYED WITH THIS PRODUCT. EXTREME CARE MUST BE TAKEN TO AVOID CONTACT WITH CROPS OR DESIRABLE PLANTS WITHOUT THE ROUNDUP READY\* TECHNOLOGY, OR WITH NATIVE VEGETATION, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.



**Section 7. COTTON - ROUNDUP READY FLEX\*  
FOR OVER-THE-TOP (OTT) APPLICATIONS MADE IN ROUNDUP READY FLEX\* COTTON FROM CROP EMERGENCE TO HARVEST**

No more than 4 applications may be made OVER THE TOP in any one crop. Any single application MUST NOT exceed 2.2 L/ha. Applications MUST NOT be made between 22 NODES and 60% BOLL OPEN STAGE.	One (1) of the four (4) applications may be made OVER THE TOP in any one crop between 60% BOLL OPEN STAGE and HARVEST. Application at this stage MUST NOT exceed 2.2 L/ha. NO MORE THAN FOUR (4) APPLICATIONS MAY BE MADE IN ANY ONE CROP.
---	--

Tank-mixtures with other herbicides or insecticides are not recommended for over-the-top applications of this product due to the potential for reduced weed control or crop injury to result (except Bouncer® 960S Herbicide).

Tank mixes with Dropp® may be used providing the crop is 60% open and immature bolls cannot be cut with a sharp knife, alternatively where the seed coat in bisected bolls is black in colour.

No additional surfactant is required for use in Roundup Ready Flex\* Cotton.

NOTE: The total glyphosate active constituent applied in any one crop must not exceed 4.14 kg/ha (including pre-harvest application).

**SITUATION** – Cotton with Roundup Ready Flex\* Technology

OTT UP TO 22 NODES (except when mixed with Bouncer® 960S Herbicide where application is between 4 and 18 nodes only)

NO MORE than FOUR (4) applications are permitted in crop up to 22 nodes.

Any single application in crop up to 22 nodes MUST NOT exceed 2.2 L/ha.

WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Afghan melon, African turnip weed, Annual ryegrass, Annual ground cherry, Barnyard grass, Bathurst burr, Black pigweed, Bladder ketmia, Boggabri weed, Button grass, Caltrop, Camel melon, Caustic weed, Columbus grass, Datura, Dead nettle, Liverseed grass, Lovegrass, Mexican poppy, Milk thistle, Mintweed, Native millet, New Zealand spinach, Noogoora burr, Paradoxa grass, Pigweed (up to 25cm diam.), Sow thistle, Spear thistle, Stinkgrass, Sweet Summer grass, Thornapple, Turnip weed, Urochloa, Variegated thistle, Volunteer cereals, Volunteer sorghum, Wild oats, Wild/Prickly lettuce, Wireweed, Yellow vine	0.77-2.2 L/ha	Rate Selection Use the lower rates on young weeds and increase to the higher rate where weeds are dense or well developed. Dense infestations of some weeds e.g. Barnyard grass, Liverseed (Urochloa) grass may need follow up treatments for complete control.
Climbing buckwheat (less than 12 leaves), Couch, Johnson grass	1.4-2.2 L/ha	Use the higher rate on plants at the flowering/seed head stage. For Johnson grass apply to plants with a minimum of 30 cm new growth. For long term control of Couch and Johnson grass, repeat applications will be required.
Nutgrass	2.2 L/ha followed by 2.2 L/ha	Make first application to actively growing plants when the majority of nutgrass plants have reached at least the 6-8 leaf stage but preferably later. Allow for maximum re-emergence before retreating.
<u>The following weeds are controlled pre-emergence by Bouncer® 960S Herbicide, this tank mix will also control all weeds listed in Section 2 with minimum application rates of 1.5-2.2 L/ha Nufarm weedmaster DST:</u> Awless barnyard grass, Barnyard grass, Dinebra ( <i>Dinebra retroflexa</i> ), Crowfoot grass, Liverseed grass, Lovegrasses, Pigeon grasses ( <i>Setaria spp.</i> ), Summer grass, Wandering Jew ( <i>Tradescantia fluminensis</i> )	1.5-2.2 L/ha + 1 L/ha Bouncer® 960S Herbicide	<b>Over the top (OTT) application:</b> Apply from the 4 node to 18 node crop growth stage only. Apply when weeds are present, using a boom sprayer applying 85-120 L of mixture per hectare. For maximum efficacy ensure even distribution of spray across the furrow. Some minor transient leaf spotting or leaf scorching may occur after application. To avoid any leaf spotting or leaf scorching use a directed or shielded spray. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4cm of soil. If rain does not occur or irrigation is unavailable to incorporate reduced residual control may occur. Apply only one application of Bouncer 960S herbicide per crop. <b>DO NOT</b> use both an OTT and directed spray application (see below).

**SITUATION** – Cotton with Roundup Ready Flex® Technology

Apply between 60% BOLL OPEN STAGE and HARVEST

NOT MORE than one (1) Application.

**DO NOT** use on crops intended for seed production

Application made between 60% open stage and harvest MUST NOT exceed 2.2 L/ha.

WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Bathurst burr, Noogoora burr, Winter annual weeds including Milkthistle, Common sowthistle	1.0-2.2 L/ha	Use the lower rate on light infestations of small weeds, where the crop canopy allows adequate spray coverage of the weeds. Increase to the higher rate when the crop canopy may limit spray coverage, when treating dense infestations, or when treating larger weeds. Apply alone or in tank mixtures with Dropp®. Apply when at least 60% of bolls are open and immature bolls cannot be easily cut with a sharp knife. Alternatively, where the seed coat in bisected bolls is black in colour. Where a leafy canopy limits spray coverage, reduced weed control can be expected. For best results under these conditions, delay application until canopy re-opens following initial conditioning treatment.

**FOR DIRECTED SPRAYING IN TANK-MIXES IN ROUNDUP READY FLEX® COTTON**

NOTE: The restrictions for OTT spraying regarding the number of applications and the total amount of glyphosate active constituent do not apply to this use pattern. This use can be in addition to the OTT uses listed previously in this section of the label.

SITUATION & CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Roundup Ready Flex cotton (4 node–18 node)	<u>The following weeds are controlled pre-emergence by Rifle® 440 Herbicide, this tank mix will also control all weeds listed in Section 2 with minimum application rates of 1.2-1.7 L/ha Nufarm weedmaster DST:</u> Barnyard grass, Button grass, Common heliotrope ( <i>Heliotropium europeum</i> ), Crested goosefoot ( <i>Chenopodium cristatum</i> ), Crowsfoot grass ( <i>Eleusine indica</i> ), Dwarf amaranth ( <i>Amaranthus macrocarpus</i> ), Early Spring grass ( <i>Eriochloa</i> spp.), Green amaranth ( <i>Amaranthus viridis</i> ), Liverseed grass, Mexican clover ( <i>Richardia brasiliensis</i> ), Mossman River grass ( <i>Cenchrus echinatus</i> ), Native millet, Pale pigeon grass ( <i>Setaria glauca</i> ), Paspalidium ( <i>Paspalidium</i> spp.), Pepper grass ( <i>Panicum whitei</i> ), Pigweed, Queensland bluegrass ( <i>Dichanthium sericeum</i> ), Red Flinders grass ( <i>Iseilema vaginiflorum</i> ), Scarlet pimpernel ( <i>Anagallis arvensis</i> ), Small burr grass ( <i>Tragus australianus</i> ), Stinkgrass, Weeping lovegrass ( <i>Eragrostis parviflora</i> ), Wireweed. <u>The following weeds are suppressed pre-emergence by Rifle® 440 Herbicide:</u> Caltrop, Common sowthistle, Common verbena ( <i>Verbena officinalis</i> ), Mintweed, Pepper cress ( <i>Lepidium</i> spp.), Stagger weed ( <i>Stachys arvensis</i> )	1.5-2.2 L/ha + 3.4 L/ha Rifle® 440 Herbicide	<b>Directed spray applications only:</b> Apply after weeds have emerged as a directed spray application to avoid shading by the crop which may affect herbicide placement. For maximum efficacy ensure even distribution of spray across the furrow. Rainfall or irrigation must occur within 10 days of application for product incorporation into the soil. <b>DO NOT</b> apply after full flowering (BBCH 65). Leaf yellowing and crinkling will usually occur after application. Apply only one application of Rifle 440 Herbicide per crop.
Roundup Ready Flex cotton (4 node–18 node), cont.	<u>The following weeds are controlled pre-emergence by Bouncer® 960S Herbicide:</u> Awless barnyard grass, Barnyard grass, Dinebra, Crowsfoot grass, Liverseed grass, Lovegrasses, Pigeon grasses, Summer grass, Wandering Jew	1.5-2.2 L/ha + 1 L/ha Bouncer® 960S Herbicide	<b>Directed spray applications only:</b> Apply after weeds have emerged as a directed spray application to avoid shading by the crop which may affect herbicide placement. For maximum efficacy ensure even distribution of spray across the furrow. Some minor transient leaf spotting or leaf scorching may occur after application. To avoid any leaf spotting or leaf scorching use a directed or shielded spray. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4cm of soil. If rain does not occur or irrigation is unavailable to incorporate reduced residual control may occur. Apply only one application of Bouncer 960S herbicide per crop. <b>DO NOT</b> use both an OTT and directed spray application (see above).

**WARNING:** THE APPLICATIONS RECOMMENDED ABOVE ARE FOR USE ONLY WITH IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE ROUNDUP READY FLEX® TECHNOLOGY. SEVERE INJURY OR DEATH OF COTTON WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY FLEX® TECHNOLOGY ARE SPRAYED WITH THIS PRODUCT. EXTREME CARE MUST BE TAKEN TO AVOID CONTACT WITH CROPS OR DESIRABLE PLANTS WITHOUT THE ROUNDUP READY® TECHNOLOGY, OR WITH NATIVE VEGETATION, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

**Section 8. PRE AND POST HARVEST USES, PASTURE TOPPING**

**GENERAL COMMENT**

**DO NOT** use on crops intended for seed production or sprouting.

**DO NOT** apply more than one (1) pre-harvest application per crop.

**DO NOT** apply if heavy rains are imminent.

**DO NOT** apply to malting barley.

Apply with ground boom or aerial equipment. For pre-harvest/cutting applications, speed of crop desiccation is dependent on crop stage, growing conditions and weather conditions during and after application. Any subsequent weed management strategies should involve an integrated weed management (IWM) approach to minimize development of glyphosate resistance.

CROP/SITUATION	SITUATION / WEEDS	RATE	CRITICAL COMMENTS
<b>PRE-HARVEST/CUTTING APPLICATIONS</b>			
BARLEY (except malting barley)	As a harvest aid Annual weeds	1.9 L/ha	Apply to mature crop from late dough stage (28% moisture) onwards. <b>DO NOT</b> harvest within 7 days after application.
CANOLA (including conventional, triazine tolerant, CLEARFIELD*, TruFlex* and Roundup Ready* varieties)  MUSTARD (oilseed cultivars) ( <i>Brassica juncea</i> )	As a harvest aid Annual weeds	1.4-4.1 L/ha	<b>DO NOT</b> use as a pre-harvest/cutting application on canola hybrids with the Optimum GLY* herbicide tolerance trait. Application on canola hybrids with the Optimum GLY* herbicide tolerance trait can only at or before BBCH 61 (refer to section 4 for further information). Apply to mature standing crop from early senescence (minimum of 20% of canola/mustard seeds as a random visual sample from various heights in the crop canopy from the main stem have changed to a dark brown/black colour) prior to windrowing or direct harvest. For further information on timing contact your Nufarm representative. <b>Application can also be made at the time of windrowing (windrow equipment fitted with spray booms). To avoid shatter losses from ground boom application; apply before complete senescence of the crop.</b> Use the higher rate when crops or weeds are dense and where faster desiccation is required. <b>DO NOT harvest for 5 days after application to standing crops.</b> <b>DO NOT apply after completion of the windrowing process</b> <b>DO NOT overspray windrows.</b> <b>DO NOT direct spray at windrows</b> <b>DO NOT apply to standing crops and again at the time of windrowing.</b> For application to standing crops a minimum water rate of 80 L/ha is recommended to ensure adequate coverage of target weeds below the crop canopy.
COTTON (conventional)	Bathurst burr Noogoora burr Winter annual weeds including: Common sowthistle /milkthistle	1.0-1.9 L/ha	Use the lower rate on light infestations of small weeds, where the crop canopy allows adequate spray coverage of the weeds. Increase to the higher rate when the crop canopy may limit spray coverage, when treating dense infestations, or when treating larger weeds. Apply alone or in tank mixtures with Dropp*. Apply when at least 60% of bolls are open and immature bolls cannot be easily cut with a sharp knife. Where a leafy canopy limits spray coverage, reduced weed control can be expected. For best results under these conditions, delay application until canopy re-opens following initial conditioning treatment.
	Nutgrass (seasonal suppression only)	1.9 L/ha	Where control of Nutgrass or Noogoora burr is required treatments should be applied prior to the onset of frosts. When tank mixed with defoliant, a slightly higher proportion of cotton leaf may be retained, particularly where the higher rate is used. Read and follow all label direction for the tank mix products. <b>For pre-harvest application in Roundup Ready Flex* Cotton, please refer to Section 7.</b>
HAY/SILAGE production from annual pastures or crops	Prevention of re-growth Annual weeds	1.4-4.1 L/ha	Apply to mature standing pasture within 1–10 days prior to cutting or mowing. Annual pasture includes oats, wheat, triticale, barley, annual ryegrass and other annual forage grasses cut for hay or silage use. Annual pasture also includes broadleaf crops such as canola, pulses, etc., and forage brassicas, etc. cut for hay or silage use. Refer to uses in this table for crops where pre-harvest application is allowed for grain production. Use the higher rate if the hay/silage is of high density, if cutting is planned within 3 days of application, or if the crop is rank or lodged and where faster desiccation is required. The lower rate is for low density hay and silage pasture only. <b>DO NOT</b> cut within 1 day after application (except for crops specifically listed elsewhere in this table where longer withholding periods are required). A minimum of 3 days prior to cutting is recommended where conditions result in slow translocation in the target plant. Good spray coverage is also important for best results and it is recommended that a minimum application water volume of 70 L/ha is used.

CROP/SITUATION	SITUATION / WEEDS	RATE	CRITICAL COMMENTS
PULSE CROPS, including: Adzuki beans Chickpeas Cowpea Faba beans Field peas Lentils Mungbeans Soybean Vetch	Crop desiccation As a harvest aid Annual Weeds	0.77-2.0 L/ha	Use higher rates where crops or weeds are dense and where faster desiccation is required. Application should be made at or after crop maturity: <b>Chickpeas and Lentils</b> - apply when physiologically mature and less than 15% green pods. <b>Faba beans</b> - apply when pods turn black and average seed moisture content is below 30%. <b>Field peas</b> - apply when seeds turn yellow and average seed moisture content is below 30%. <b>Mungbeans / Adzuki beans and Cowpea</b> - apply to mature crops when pods are brown/black. <b>Soybean</b> - apply only after seed pods have lost all green colour and 80-90% of leaves have dropped. <b>Vetch</b> - apply at or after crop maturity when seed moisture content is below 30%. <b>DO NOT harvest within 7 days of application.</b>
		<b>Chickpeas only:</b> 0.6-1.2 L/ha plus 5 g/ha Associate® Herbicide	Apply when chickpeas are physiologically mature and less than 15% of green pods are present. Use higher rates where crops or weeds are dense and where faster desiccation is required. <b>DO NOT harvest within 7 days of application.</b>
SORGHUM	As a harvest aid Annual weeds and sorghum ratoon growth	1.15-2.3 L/ha	<b>DO NOT</b> apply if crop is under stress from low moisture, frost, cold or waterlogging. <b>RATE SELECTION</b> Use the lower rate for control of crop and late tillers and suppression of ratoon regrowth. Use the higher rate for improved suppression of ratoon regrowth, where the crop has produced significant number of late tillers or where following crops will be established without further treatment. <b>TIMING</b> Apply when grain moisture is less than 25%. Application can be made when moderate browning has occurred. <b>CAUTION</b> Treatment may increase potential for CROP LODGING, particularly if prior moisture stress has occurred. Under any set of environmental conditions, individual varieties can vary in response to preharvest treatments. In general, varieties with a more "determinant" growth habit are more susceptible than "indeterminant" varieties. Harvest should commence at least 7 days after application provided sufficient dry down has occurred to avoid possible lodging. <b>CAUTION Sorghum may be naturally toxic to stock.</b>
WHEAT	As a harvest aid Annual weeds	1.4-4.1 L/ha	Apply to mature crop from late dough stage (28% moisture) onwards. The higher rate will be required when crops are heavy and leaf shading effects may occur. <b>DO NOT harvest within 5 days after application.</b>
<b>POST-HARVEST APPLICATIONS</b>			
SORGHUM Harvested crops	Sorghum regrowth	0.77-1.15 L/ha for fresh regrowth from slashed stubble	<b>APPLY UNDER GOOD GROWING CONDITIONS ONLY. DO NOT</b> apply if plants are under stress from low moisture, frost, cold or waterlogging. <b>SLASHED STUBBLE AND SPRING REGROWTH</b> Apply when fresh regrowth is at least 20cm high. <b>STANDING STUBBLE</b> Apply only if sufficient green leaf is present. If grazing has occurred allow regrowth to 20cm before treatment. <b>RATE SELECTION</b> Use the lower rate for knockdown and regrowth suppression where cultivation is to follow. Increase to the higher rate for improved regrowth control. <b>NOTE</b> Variable results occur where the crop has been subject to stress or growing conditions are marginal. <b>CAUTION Sorghum may be naturally toxic to stock.</b>
		1.15-1.5 L/ha for standing stubble if sufficiently green and for fresh Spring regrowth	
SUGARCANE Harvested crops	Sugarcane ratoon regrowth	4.6-6.9 L/ha	<b>APPLY UNDER GOOD GROWING CONDITIONS ONLY</b> to actively growing ratoons 60-120 cm tall. <b>DO NOT</b> apply if plants are under stress from low moisture or waterlogging. Use the lower rate for suppression or where cultivation is to follow. Use higher rate for control.
<b>PASTURE TOPPING</b>			
PASTURES (For annual and perennial grasses, Bent grasses, Serrated Tussock, Capeweed and Calomba daisy seed-set reduction)	Annual ryegrass	350 mL/ha	Reduction in pasture legume population may occur because of treatment. Remove stock prior to treatment to allow even regrowth. Apply to Capeweed and Annual ryegrass at FLOWERING. For other grasses, apply from HEAD to MILKY DOUGH stage. Use the higher rate for dense infestations or where Annual ryegrass is present. Apply before signs of plants "haying off".
	Barley grass	230-350 mL/ha	
	Brome grass	230-350 mL/ha	
	Calomba daisy	350 mL/ha	
	Capeweed	230-350 mL/ha	
	Silvergrass	230-350 mL/ha	
Bent grasses	285-480 mL/ha	<b>TIMING</b> Treat from late October to late November. Apply before seed heads have emerged. Use the higher rate where growth is excessive and renovation is intended the following Autumn. <b>FOLLOW-UP MANAGEMENT</b> Graze hard after spraying.	
Serrated Tussock	600-1000 mL/ha	Apply to actively growing and stress free plants. Best results obtained during mid September – mid October. Apply prior to any seed head emergence. Also see Aerial Equipment. Surfactants: Addition of 200 mL of Wetter TX™ to 100 L of spraying solution may improve results. Rates: The lower rates will be less damaging to desirable pasture species. If seed head emergence is imminent then higher rates will give better results.	

**Section 9. OPTICAL SPOT SPRAY TECHNOLOGIES**

Note: Calibrate the sprayer to spray the equivalent of 100 L/ha

For weed cover between 0 and 30 %. If percentage weed cover exceeds 30 % use approved boom spray rates.

SITUATION	WEEDS	RATE	CRITICAL COMMENTS
FALLOW	Australian bindweed	3.5-7.0 L/100L	Suppression only at low rates. Use the high rate on larger mature plants
	Barnyard grass		Use the high rate on large mature plants
	Bladder ketmia		Apply low rate to rosette to flowering plants. Use the high rate on large mature plants
	Common sowthistle		
	Turnip weed		
	Yellow vine (caltrop)		
Prior to sowing the following winter or summer broadacre crops, or starting a fallow and for fallow maintenance: Cereals; Pulses; Canola; Mustard (oilseed cultivars) ( <i>Brassica juncea</i> ); Cotton; as per the plantback table on the Terra'd or® 700WG Herbicide product label	Annual grass and broadleaf weeds as listed in <b>Section 2 weed table.</b>	2.3 L/100L + 40 g/100L Terra'd or® 700WG Herbicide + 1L/100L Banjo® Spray Adjuvant*	Observe the plant-back periods listed in the GENERAL INSTRUCTIONS. Sow following crops with a seeder that will move treated soil away from crop row (e.g. knife point with press wheels). Use of seeding machinery, or planting under conditions where product or treated soil remains or moves back into the crop row may result in crop damage. This may be caused by factors such as inappropriate seeding machinery, narrow row spacings, fast travel speeds, soil type, wind, heavy rainfall or irrigation after planting, etc. This is particularly important for susceptible crops such as canola. <b>DO NOT</b> apply post-sowing pre-emergent. Addition of Banjo spray adjuvant* at 1.0 % v/v must always be used with Terra'd or® 700 WG. Rainfast after 1 hour of daylight after application to seeding annual weeds. * or other registered high quality methylated seed oil adjuvant

**Section 10. POST SOW PRE-EMERGENT (PSPE) USES, SHIELDED SPRAYERS AND BAND SPRAYING**

SITUATION & CROP	WEEDS	RATE	CRITICAL COMMENTS
COTTON (inter-row shielded sprayers)	Refer to weeds and corresponding application rates in Section 2.		Apply Nufarm weedmaster DST to weeds growing between crop rows using a shielded sprayer. <b>DO NOT</b> apply in crops less than 20 cm high. <b>DO NOT</b> allow spray or spray drift to contact any part of the cotton plant as severe injury or destruction may result. Can be used in conventional and tolerant cotton varieties.
ONIONS (PSPE)	For control of annual weeds and suppression of perennial weeds, including Rope Twitch	0.77-2.3 L/ha	Apply post-sowing and at least 7 days before crop is due to emerge. <b>DO NOT</b> apply to emerging Onion plants as severe injury will result. Use the lower rate on small, actively growing annual weeds. Increase to the higher rate for larger annual weeds (over 15 cm tall) and for suppression of perennial weeds.
PASTURES (Band spraying) For suppression or control of pastures species prior to drilling improved pasture, forage species, Soybeans or Leucaena.	Refer to weeds and corresponding application rates in Section 2.		Apply low rates for suppression. Apply higher rates where complete control is required. Band spraying may be done immediately after the sowing operation. Mount the nozzles behind the coultter/tyne/press wheel assembly of the band seeder. Adjust to spray 0.5 to 1.0m strips. Ensure minimal disturbance of the pasture. Excessive dust created in the seeding operation may reduce herbicide activity. Pasture seed must be drilled at the appropriate depth and covered by soil. LEUCAENA: Apply maximum of 1.6 L/ha through a single taper fan nozzle LF-80 mounted at the rear of the single row planter providing a 1 m swath. Planting rows to be 4 m apart.
SUGARCANE (inter-row shielded sprayers)	Annual and perennial grasses and broadleaf weeds	1.4-5.7 L/ha	Apply to weeds growing between crop rows using a ground based hooded and shielded sprayer. Apply at early growth stage of crop, before formation of the cane. Apply no more than 3 applications, to a maximum of 13.8 L/ha per crop. <b>DO NOT</b> allow spray or spray drift to contact any part of the crop as severe injury may result.

**Section 11. BRUSH AND WOODY WEEDS – MIXTURES WITH ASSOCIATE**

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
COMMERCIAL AND INDUSTRIAL AREAS DOMESTIC AND PUBLIC SERVICE AREAS FORESTS PASTURES RIGHTS OF WAY	Blackberry ( <i>Rubus</i> spp.) Volunteer Pine ( <i>Pinus</i> spp.) wildlings (suppression only)	<b>Handgun or Knapsack</b> 320 mL Nufarm weedmaster DST plus 3 g Associate* per 100 L of water	For Blackberries, apply from flowering until prior to leaf yellowing. Due to widespread picking of Blackberries by the public, it is not recommended that the product be applied to bushes bearing mature fruit. Application to Pine wildlings less than 50cm in height should be controlled when actively growing. Use Pulse® Penetrant at the rate of 200-500 mL per 100 L water.
	Bracken ( <i>Pteridium esculentum</i> )	<b>Aerial or Boom</b> For Blackberry, Coppice and Volunteer Pine wildlings: 6.1 L/ha plus 60 g Associate* per ha For other weeds: 3.1 L/ha plus 30 g Associate* per ha	For Bracken, apply when fronds are fully unfurled but prior to first frosts. For boom application, refer to Boom application section. Use Pulse® Penetrant at the rate of 200-500 mL per 100L of water.
	Coppice ( <i>Eucalyptus globulus</i> )		Use Pulse Penetrant at the rate of 500 mL per 100 L water. For coppice control, apply using a minimum 400 L/ha using a double pass application method to coppice up to a maximum 1.5 m in height. Target every stem separately. <b>DO NOT</b> disturb for 6 weeks to allow maximum translocation.
	Gorse ( <i>Ulex europaeus</i> )		For Gorse, apply when actively growing at any time of year, except Spring. Use Pulse® Penetrant at the rate of 200-500 mL per 100 L of water.
	Lantana ( <i>Lantana camara</i> )		For Lantana, apply when actively growing. <b>DO NOT</b> apply during periods of Summer drought stress. Use Pulse® Penetrant at the rate of 200-500 mL per 100 L of water.
	St John's Wort ( <i>Hypericum perforatum</i> )		For St John's Wort, apply when actively growing from Spring to Summer. Use Pulse® Penetrant at the rate of 200-500 mL per 100 L of water.
Sweet Briar ( <i>Rosa rubiginosa</i> )		For Sweet Briar, apply when in full leaf, prior to leaf fall. Use Pulse® Penetrant at the rate of 200-500 mL per 100 L of water.	

**Section 12. UNWANTED TREE CONTROL**

Ensure trees are actively growing at time of treatment and not under stress of drought, waterlogging or cold.

METHOD	SPECIES CONTROLLED	TREE SIZE	MIXTURE (BY VOL.) Product:Water	CRITICAL COMMENTS
CUT STUMP	Jarrah ( <i>Eucalyptus marginata</i> ), Long-leaved box ( <i>E. goniocalyx</i> ), Marri ( <i>E. calophylla</i> ), Messmate ( <i>E. obliqua</i> ), Narrow-leaved peppermint ( <i>E. radiata</i> )	0-10 cm basal diameter	1:15	Dilute with water in the recommended ratio. Cut tree close to ground and immediately wet stump surface thoroughly using Splatter Gun, spray, swab or bush. Remove any branches on the stump and treat any cut surface.
	Privet ( <i>Ligustrum</i> spp.), Rhus ( <i>Toxicodendron succedaneum</i> )	0-30 cm basal diameter	1:1	
FOLIAR APPLICATION: Low Volume (Gas Gun or Splatter Gun)	Bullich ( <i>Eucalyptus megacarpa</i> ), Marri, Jarrah	0-1.5 m height	1:15 Add Pulse* Penetrant at 20 mL/10L spray mixture	Dilute in the recommended ratio. Calibrate Splatter Gun to apply 5 mL of solution per dose, as a fine spray. Apply 5 mL per 0.5 m tree height. Ensure spray contacts all foliage.
	<i>Eucalyptus</i> spp.	0-1.5 m height	1.5 Add Pulse* Penetrant at 20 mL/10 L spray mixture	
FOLIAR APPLICATION: High Volume (Knapsack or Handgun)	<i>Eucalyptus</i> spp., Willows ( <i>Salix</i> spp.)	0-2 m height	0.8-1.0 L/100 L For <i>Eucalyptus</i> spp. Add Pulse* Penetrant at 200 mL/100 L spray mixture	Spray to wet all foliage. Use the higher rate for trees 1 to 2 m high.
STEM INJECTION	Flooded gum ( <i>Eucalyptus grandis</i> ), Ghost gum ( <i>Corymbia papuana</i> ), Gum-topped bloodwood ( <i>C. dichromophloia</i> ), Messmate, Narrow-leaved ironbark ( <i>E. crebra</i> ), Pink bloodwood ( <i>C. intermedia</i> ), Poplar box ( <i>E. populnea</i> ), Privet, Rhus, Silver-leaved ironbark ( <i>E. melanophloia</i> ), Silvertop ash ( <i>E. sieberi</i> ), Spotted gum ( <i>C. maculata</i> ), Swamp mahogany ( <i>Lophostemon suaveolens</i> ), White mahogany ( <i>E. acmenoides</i> ), Willows	0-25 cm basal diameter	Undiluted 1 mL/cut	Use an applicator calibrated to deliver 1 or 2 mL per cut. Make 5 cm cuts at an oblique angle to penetrate the sapwood beneath the bark. Space cuts at 13 cm centers around tree circumference below any branching, otherwise remove or treat all branches below cuts. On multiple trunk trees ensure each trunk is treated.
	Camphor laurel ( <i>Cinnamomum camphora</i> )	Over 25-60 cm basal diameter	Undiluted 2 mL/cut	
		Basal diameter to 25 cm	Mixture 1:1 2 mL/cut	
	Basal diameter over 25-60 cm	Undiluted 2 mL/cut		

**Section 13. GENERAL WEED CONTROL**

SITUATION	CRITICAL COMMENTS
<p><b>GENERAL WEED CONTROL</b> In Domestic Areas (Home Gardens), Commercial, Industrial and Public Service Areas, Agricultural Buildings and Other Farm Situations.</p>	<p>Refer to Section 2 for boom spray rates or Section 14 for handgun or knapsack rates.</p> <p>For the control of many grasses and broadleaf weeds. <b>RATE: 8 mL per litre of water.</b> When treating dense, maturing weed populations in non-arable scenarios using a boom sprayer, ensure good coverage by applying Nufarm weedmaster DST at a rate of no more than 6 mL per litre of water. Apply when weeds are actively growing. Apply to ensure complete and uniform wetting of foliage. Visible symptoms may take from 3 to 7 days to develop.</p>
<p><b>AGRICULTURAL AREAS</b></p>	<p>Nufarm weedmaster DST may be used for control of annual, perennial and woody weeds as directed, in agricultural land prior to sowing of any edible or non-edible crop, but not prior to transplanting tomato seedlings.</p>
<p><b>DRY DRAINS AND CHANNELS ONLY</b></p>	<p><b>DO NOT</b> apply to weeds growing in or over water. <b>DO NOT</b> spray across open bodies of water, and <b>DO NOT</b> allow spray to enter the water. <b>DO NOT</b> allow water to return to dry channels and drains within 4 days of application.</p>
<p><b>FORESTS</b></p>	<p>Nufarm weedmaster DST may be used prior to establishment of nurseries, for site preparation prior to planting and amongst established trees using a directed or shielded spray, or using selective wiper equipment. <b>DO NOT</b> allow wiper surface to contact any part of the tree. <b>DO NOT</b> allow spray or spray drift to contact foliage or green bark of desirable trees, since severe injury may result. For improved control of woody and herbaceous weeds pre-plant (broadcast spraying by air or helicopter) or post-plant (inter-row only), add 0.5-1.5 L/ha of Comet 400 with a rate of Nufarm weedmaster DST up to 6.1 L/ha</p>
<p><b>NON-AGRICULTURAL AREAS</b> Around Buildings, Commercial and Industrial Areas, Domestic and Public Service Areas, Right-Of-Ways.</p>	<p>Nufarm weedmaster DST does not provide residual weed control. For residual control of annual weeds, Nufarm weedmaster DST may be tank mixed with certain residual herbicides. See Tank Mixtures/Compatibility.</p>
<p><b>PASTURE</b></p>	<p><b>DIRECTED (SPOT) APPLICATION</b> Nufarm weedmaster DST is non-selective and may damage or kill any plant in the sprayed area. Re-treatment and/or pasture improvement may be necessary to restrict seedling re-establishment. <b>SELECTIVE APPLICATION</b> See Wiper Equipment. <b>BOOM APPLICATION</b> Nufarm weedmaster DST may be used to suppress or kill existing pasture species prior to re-seeding or establishment of other crops. When spot application (spray or wiper) is undertaken, grazing stock need not be removed. <b>CAUTION</b> Certain plants may be naturally toxic to stock. Where known toxic plants are present <b>DO NOT</b> allow stock to graze until complete browning of treated plants has occurred.</p>
<p><b>TREE AND VINE CROPS</b> Vineyards, Berries and Other Small fruits (Excluding Strawberry), Citrus fruits, Tropical and Sub-Tropical fruits, Pome fruits, Stone fruits, Tree nuts, Duboisia, Hops, Tea</p>	<p>Apply as a directed or shielded spray or using wiper equipment. <b>DO NOT</b> apply as spray near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift. <b>DO NOT</b> allow wiper surface to contact any part of the tree, vine or plant. <b>Citrus fruit, Nuts, Olives, Pome fruit &amp; Vineyards DO NOT</b> allow spray or spray drift to contact green bark or stems, canes, laterals, suckers, fresh wounds, foliage or fruit. <b>Hops</b> Apply in Winter, prior to crop emerging from dormancy. <b>Tea</b> Apply a maximum of 3.1 L/ha by shielded boom or directed off-centre nozzle or 380 mL/100 L by directed hand-gun or knapsack to avoid application to the crop. <b>All other crops DO NOT</b> allow spray or spray drift to contact any part of the plant including the trunk. <b>CAUTION</b> Where split bark on Kiwifruit and green stems on Pawpaw occur, extreme care is required.</p>
<p><b>WIPER APPLICATION</b> (above a crop or pasture)</p>	<p>Wiper equipment (e.g. Ropewick, canvas, felt or carpet applicators) may be used to apply Nufarm weedmaster DST. Avoid contact with desirable vegetation. Operate wiper equipment a minimum of 10 cm above the crop or pasture. Weeds should be at least 15 cm above the crop or pasture at time of application. Speed of travel should be no greater than 8 km/h. Best results are achieved at lower speeds and where two applications are made in opposite directions (double pass). Where weeds are of variable height, or occur in dense infestations or clumps, some plants may not be contacted by the herbicide solution. In these cases repeat treatment may be necessary. <b>RATE:</b> Mix 770 mL Nufarm weedmaster DST with 2.2 L clean water. Adjust flow rate to suit equipment.</p>

## Section 14. WEED CONTROL TABLE FOR SECTION 13 (HANDGUN, KNAPSACK ONLY)

WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
<b>ANNUAL WEEDS</b> Amaranth, Bathurst burr, Barley grass, Billygoat weed, Brome grass, Barnyard grass, Caltrop, Canary grass, Capeweed, Chickweed, Cobblers peg, Deadnettle, Doublegee, Fumitory, Ground cherry, Hedge mustard, Lesser swinecress, Liverseed grass, Mintweed, Noogoora burr, Paradoxa grass, Paterson's curse, Pigweed, Potato weed, Ryegrass, Saffron thistle, Silvergrass, Sow thistle, Spear thistle, Spiny burgrass, Spurge, Sub clover, Thornapple, Wild mustard, Wild oats, Wild turnip, Winter grass, Variegated thistle, Volunteer cereal	<b>Handgun:</b> 375-550 mL/100L <b>Knapsack:</b> 55-80 mL/15L <b>Boom:</b> Refer to Section 2	Use higher rate on weeds over 15 cm in height or diameter or where dense weed cover limits spray coverage. Use higher spot spraying rate when applying less than 5 L spray per 100sqm. Nufarm weedmaster DST does not provide residual weed control. Repeat treatments may be necessary to control later germinating weeds. For residual control of annual weeds Nufarm weedmaster DST may be tank-mixed with certain residual herbicides. See Tank Mixtures in the General Instructions for directions. <b>DO NOT</b> use an atrazine tank-mix for control of barnyard grass or liverseed grass.
<b>PERENNIAL WEEDS</b> Artichoke thistle, African lovegrass, Bent grass, Carpet grass, Cocksfoot, Flatweed, Johnson grass, Kangaroo grass, Kikuyu, Navua sedge, Nutgrass, Paspalum, Phalaris, Plantains, Poa tussock, Prairie grass, Qld blue grass, Red-leg grass, Rhodes grass, Rope twitch, Sorrel, Soursob, Yorkshire fog	<b>Handgun:</b> 540-760 mL/100L <b>Knapsack:</b> 80-115 mL/15L <b>Boom:</b> Refer to Section 2	Control of established perennials is best obtained when plants are at the seedhead stage. In general, best control of Winter growing perennials is obtained with application during Winter-Spring. Best control of Summer growing perennials is obtained with application late Summer and Autumn. For Nutgrass in cultivated situations apply sequential low rate treatments when Nutgrass has a minimum of 6-8 leaves. Use the higher rate in uncultivated situations. For Rhodes grass, Rope twitch, Prairie grass, Qld blue grass, Johnson grass, Kangaroo grass, Kikuyu, Redleg grass, Paspalum and Sorrel, use the higher rates only.
Blady grass, Bracken, Couch, Guinea grass *Paragrass, Silverleaf nightshade, *Water couch *Use on Dry Drains and Channels ONLY (see Use Situations critical comments above)	<b>Handgun:</b> 1.0-1.54 L/100L <b>Knapsack:</b> 150-230 mL/15L <b>Boom:</b> Refer to Section 2	For Bracken add Pulse® at 200-500 mL/100L spray mix. Best control of couch in WA and SA is obtained with Spring treatment. Most effective control of couch in eastern states is obtained with Summer and Autumn treatments. In cultivated situations use sequential treatments of 2.3-4.8 L/ha for control. Only use higher rate for handgun and knapsack for Silverleaf nightshade.
<b>WOODY WEEDS</b> Bamboo ( <i>Phyllostachys</i> spp), Bitou bush ( <i>Chrysanthemoides monilifera</i> subsp. <i>rotundata</i> ), Boneseed ( <i>C. monilifera</i> subsp. <i>monilifera</i> ), Boxthorn ( <i>Lycium ferocissimum</i> ), Crofton weed ( <i>Ageratina adenophora</i> ), Gorse, Groundsel bush ( <i>Baccharis halimifolia</i> ), Lantana, Mistflower ( <i>Ageratina riparia</i> )	<b>Handgun:</b> 380-770 mL/100L <b>Knapsack:</b> 60-115 mL/15L <b>Boom:</b> Refer to Section 2	Further treatment may be necessary to restrict seedling re-establishment. <b>Bamboo</b> , apply when foliage/regrowth is 1-2m tall, use higher rate only. <b>Bitou bush/Boneseed</b> , apply higher rate on bushes greater than 1.5m. Best results are achieved when treated at peak flower during Winter. <b>Boxthorn</b> minimum rate is 600 mL/100L for handgun and 80 mL per 15 L for knapsack. <b>Groundsel bush</b> , apply higher rate on bushes greater than 2 m. <b>DO NOT</b> apply in Winter. Minimum rate is 600 mL/100L for handgun and 80 mL per 15 L for knapsack. <b>Gorse, always add Pulse®</b> at 200-500 mL/100L of spray mix, use higher rate only. <b>Lantana</b> , use higher rate only. Addition of Pulse® (200-500 mL/100L) may improve control. <b>Boxthorn, Gorse, Lantana</b> Removal of bushes (after complete brownout), pasture improvement or further treatment are recommended to control seedlings and/or regrowth.
<b>WOODY WEEDS</b> Blackberry, Eucalyptus spp. (seedlings less than 2m), Hawthorn, Pampas grass ( <i>Cortaderia</i> spp.), Chinese scrub or Sifton bush ( <i>Cassinia arcuata</i> ), Sweet briar, Willow (less than 2m)	<b>Handgun:</b> 0.77-1.5 L/100L <b>Knapsack:</b> 115-160 mL/15L <b>Boom:</b> Refer to Section 2	Removal of bushes (after complete brownout), pasture improvement or further treatment are recommended to control seedlings and/or regrowth. <b>Blackberry</b> , apply from flowering to leaf fall, use higher rate on old dense infestations greater than 2 m high. In Tasmania, <b>DO NOT</b> treat bushes bearing mature fruit. <b>Chinese scrub</b> , use higher rates on bushes greater than 1m. <b>Eucalyptus spp.</b> , add Pulse® at 200-500 mL/100 L of spray mix. <b>Hawthorn</b> , apply from flowering to leaf fall, use higher rates on bushes greater than 2m. <b>Pampas grass</b> , allow regrowth to reach 1 m, best results-apply after flowering. <b>Sifton bush</b> , use higher rates on bushes greater than 1 m. <b>Sweet briar</b> , apply from late flowering to leaf fall, use 1.15-1.5 L/100L for handgun, and 160-230 mL per 15 L for knapsack, use higher rates on bushes greater than 1.5 m.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**



**WITHHOLDING PERIOD:****GRAZING:****BARLEY, CANOLA, MUSTARD:**

- DO NOT CUT OR GRAZE FOR STOCKFOOD FOR 7 DAYS AFTER APPLICATION.

**COTTON:**

- DO NOT GRAZE OR CUT FOR STOCKFOOD.

**HAY/SILAGE (OTHER THAN CROPS LISTED SEPARATELY IN THIS SECTION):**

- DO NOT MOW OR CUT FOR 1 DAY AFTER APPLICATION. A MINIMUM 3 DAYS PRIOR TO CUTTING IS RECOMMENDED WHERE CONDITIONS RESULT IN SLOW TRANSLOCATION IN THE TARGET PLANT.

**WHEAT:**

- DO NOT CUT OR GRAZE FOR STOCKFOOD FOR 5 DAYS AFTER APPLICATION.

**ALL OTHER USES:**

- NOT REQUIRED WHEN USED AS DIRECTED.

**HARVEST:****BARLEY, PULSES, SORGHUM:**

- DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

**CANOLA, MUSTARD:**

- NOT REQUIRED WHEN USED AS DIRECTED ON CANOLA HYBRIDS WITH THE OPTIMUM GLY\* HERBICIDE TOLERANCE TRAIT.
- NOT REQUIRED WHEN USED SOLELY IN-CROP OVER ROUNDUP READY CANOLA PRIOR TO BUD FORMATION.
- DO NOT HARVEST FOR 5 DAYS AFTER APPLICATION TO A STANDING CROP AS A PRE-HARVEST / CUTTING APPLICATION.
- NOT REQUIRED WHEN USED AT WINDROWING.

**COTTON**

- NOT REQUIRED WHEN USED AS DIRECTED.

**WHEAT:**

- DO NOT HARVEST FOR 5 DAYS AFTER APPLICATION.

**ALL OTHER USES:**

- NOT REQUIRED WHEN USED AS DIRECTED.

**TANK MIXTURES: REFER TO TANK MIX PARTNER LABEL AND FOLLOW ACCORDINGLY.**

**EXPORT OF TREATED PRODUCE**

Growers should note that MRL's or import tolerances **DO NOT** exist in all markets for produce treated with Nufarm weedmaster DST Herbicide. If you are growing produce for export, please check with Nufarm Australia Limited for the latest information on MRL's and import tolerance before using Nufarm weedmaster DST Herbicide.

**GENERAL INSTRUCTIONS**

Nufarm weedmaster DST is a non-volatile, non-selective, water soluble liquid herbicide with non-selective herbicidal activity. It is absorbed by plant foliage and green stems and moves through the plant from the point of contact to and into the root system. Effects may not be apparent for 3–7 days (annual weeds) or 2–3 weeks (perennial weeds) or longer under cool, cloudy conditions.

Nufarm weedmaster DST will control emerged weeds only, and provides no residual weed control. Apply treatments to weeds which have at least one true leaf (broadleaf weeds) or two leaves (grasses) to provide an adequate surface area for herbicide uptake. Nufarm weedmaster DST may be used prior to sowing any crop (edible or non-edible) but not prior to transplanting tomato seedlings.

A withholding period for grazing stock is not required in most situations (except as listed in the withholding periods section). However, it is recommended that grazing of treated plants be delayed for one day after treatment of annual weeds, or 7 days of perennial weeds are present, to ensure absorption of Nufarm weedmaster DST. Certain plants (eg. Soursob, variegated thistle) may be naturally toxic to stock. Where known toxic plants are present, **DO NOT** allow stock to graze until complete browning of treated plants has occurred.

Weeds should be actively growing at the time of treatment. **DO NOT** treat weeds under poor growing or dormant conditions (such as occur in drought, water logging, disease, insect damage or following frosts) as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust or silt. Prior herbicide application may also induce stress in weeds.

Rainfall occurring up to 6 hours after application may reduce effectiveness. Rainfastness or general efficacy may be reduced if weeds are not actively growing, are under stress or conditions of low light intensity/darkness. Heavy rainfall within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required. Delay treatment of plants wet with dew or rain, if water droplets run off when plants are disturbed. Avoid contact with foliage, green stems or fruit of crops, desirable plants and trees, since severe injury or destruction may result.

**Crop Establishment**

Nufarm weedmaster DST is recommended for control of emerged weeds prior to crop establishment. Suitable cultivation and/or sowing operations are required to provide seed bed conditions satisfactory for crop germination and development. Spraying early to control young weeds will favour preparation of suitable seed beds. On friable soils and where there is only light cover of young weeds, sowing may proceed satisfactorily from one day after spraying. In situations of heavy weed growth sowing should be delayed until weed decay and soil conditions allow formation of a satisfactory seed bed. Incorporation of green or decaying vegetation and roots into the seedbed by cultivation or sowing may cause retarded crop emergence, particularly in cold and/or wet conditions. Vegetation may be reduced by grazing and weed decay may be assisted by cultivation to leave trash on the surface. In marginal seeded conditions take care to achieve correct seeding depth, and avoid use of pre-emergence herbicides where label directions advise of risk of retarded crop emergence.

**Full soil disturbance seeding**

When cultivating or sowing with a tined implement, the operation may start one day after applying Nufarm weedmaster DST to annual species and 7 days for perennial species and should be completed within 21 days. Cultivation or sowing may start 6 hours after treatment when treating light infestations or pre-tilling annual grasses or annual broadleaved weeds less than 8cm diameter/height.

#### ***Minimal or no soil disturbance seeding***

Trash may be removed by grazing after applying Nufarm weedmaster DST. Grazing may commence one day after treatment of small annual weeds, three days for large annual weeds, and 7 days for perennial weeds, and should be completed within 21 days.

#### ***Aerial (or surface) seeding***

Delay seeding until trash is completely removed by grazing and/or plant decay. When establishing pasture, ensure application of fertilizer and insecticides and follow-up management is undertaken as required.

#### **Mixing**

Nufarm weedmaster DST mixes readily with water. Note: Reduced results may occur if water containing soil is used, eg. Water from ponds and unlined ditches, or if hard water containing calcium salts is used.

**DO NOT** mix, store or apply this product or spray solutions of this product in galvanised steel or unlined steel containers or spray tanks, since a highly flammable gas mixture may be formed. Use stainless steel, aluminium, brass, copper, fibreglass, plastic or plastic lined containers or spray tanks.

Ensure the sprayer is free of any residue of previous spray materials. Use spray solutions promptly since a gradual loss of activity will occur. Fill the spray tank with one half the required amount of clean water and add the proper amount of Nufarm weedmaster DST. Mix well before adding the remaining portion of water.

Where required, add surfactant near the end of the filling process to minimize foaming. Placing the filling hose below the surface of the spray solution will prevent excessive foaming. Removing hose from tank immediately after the filling will prevent back siphoning into water source. **DO NOT** use mechanical agitators as these may cause excessive foaming.

Spray tanks, pumps, lines and nozzles should be thoroughly rinsed with clean water following application to prevent corrosion.

#### ***Tank mixtures/compatibility***

Nufarm weedmaster DST may be tank mixed with the following herbicides, insecticides and additives. Read and follow all label directions, restraints, plant-back periods, withholding periods, regional use restrictions and safety directions for the tank mix products.

#### ***Mixing instructions for all tank mixtures***

1. Fill the spray tank 1/3 to 1/2 full with clean water and start agitation.
2. Add Nufarm Liasé where required.
3. Add recommended herbicide/insecticide/additive to the spray tank and mix thoroughly.
4. Add Nufarm weedmaster DST and the remaining water. Mix thoroughly.
5. Add surfactant, if required, near the end of the filling process to minimize foaming.
6. Always maintain adequate agitation during application and use the tank mix promptly.

#### ***Tank mixtures - Herbicide Compatibility***

Nufarm weedmaster DST is compatible with the following herbicides:

Amicide Advance<sup>®</sup> 700, Archer<sup>®</sup> 750 Dual Salt Liquid Herbicide, Associate<sup>®</sup>, Atralex<sup>®</sup>, Avadex<sup>®</sup> Xtra, Cobber<sup>®</sup> 475, Comet<sup>®</sup> 400, Nail<sup>®</sup> 240EC, Nail<sup>®</sup> 600EC, Kamba<sup>®</sup> 750 (dicamba), Estericid<sup>®</sup> Xtra 680 (2,4-D ester), Polo<sup>®</sup> LVE, Rifle 440<sup>®</sup>, Sharpen<sup>®</sup>, Simagranz, Spark<sup>®</sup>, Striker<sup>®</sup>, Terrad<sup>®</sup> or<sup>®</sup> 700WG, Terrain<sup>®</sup> 500WG, TriflurX<sup>®</sup> (trifluralin). Other brands have not been tested.

#### ***Tank mixtures - Insecticide compatibility***

Nufarm weedmaster DST is compatible with the following insecticides: Astound<sup>®</sup> Duo, Astral<sup>®</sup> 250, Chlorpyrifos 500EC, Imidan<sup>®</sup>, Kaiso<sup>®</sup> 240EG with Sorbie Technology, Le-Mat<sup>®</sup>, Lorsban<sup>®</sup> 500, Matador Zeon<sup>®</sup>, Sumithion<sup>®</sup> ULV, and emulsifiable concentrates of dimethoate and fenitrothion. Other insecticides have not been tested.

#### ***Tank mixtures - Additives***

##### **Nufarm Liasé (417 g/L ammonium sulphate liquid)**

RATE: 2 L per 100 litres spray solution.

Nufarm Liasé may be used as an adjuvant to alleviate the adverse effects of high levels of calcium, magnesium and bicarbonate ions in water. The addition of Nufarm Liasé to Nufarm weedmaster DST, when used to control annual weeds, MAY improve the performance of Nufarm weedmaster DST under adverse environmental conditions such as cool cloudy weather. Nufarm Liasé may also improve the performance of tank mixtures of Nufarm weedmaster DST and atrazine or simazine. Ammonium sulphate may be corrosive to metal parts of the sprayer. Thoroughly flush tanks, pumps and nozzles with water after use. Solubility and impurity profiles of other forms of ammonium sulphate can vary and may reduce the performance of Nufarm weedmaster DST or tank mixtures.

#### ***Surfactant addition***

Nufarm weedmaster DST requires the addition of a tank-mix surfactant, except where specifically noted in the Directions for Use.

##### **Nufarm Collide 700<sup>®</sup> Surfactant**

RATE: 250-500 mL per 100 L. The addition of Nufarm Collide 700 surfactant MAY improve weed control. At rates of 300-500 mL per 100 L, Nufarm Collide 700 is likely to modify the droplet spectrum produced by CP and flat fan nozzles. This is likely to reduce the proportion of driftable FINE droplets produced by these nozzles. For high volume application (eg. Knapsack, Handgun equipment), if Nufarm weedmaster DST is less than 1.15 L/100L, add Nufarm Collide 700<sup>®</sup> at 250 mL/100L.

##### **Nufarm Activator<sup>®</sup>**

RATE: 70-125 mL per 100 L. General Purpose non-ionic surfactants, other than Activator, may increase the production of FINE and VERY FINE droplets, which are prone to drift when used through certain nozzle types.

##### **Wetter TX Surfactant**

RATE: 200 mL/100 L spray solution. Add when treating Annual Ryegrass, Silvergrass and Perennial grasses. Wetter TX is NOT a general-purpose surfactant and should be used only where recommended. The addition of Wetter TX may improve rainfastness on Winter annual weeds.

**DO NOT** use adjuvants or surfactants other than those recommended on this label.

**DO NOT** use crop oil except when tank mixing with a herbicide for which an oil adjuvant is recommended to be used. The addition of a crop oil can reduce control of some grass weeds, particularly in Summer.

## Spray applications and spray drift risk assessment



For aerial application it is recommended where possible for this product to be applied by an aerial applicator business that holds current accreditation for the Aerial Improvement Management System –“ AIMS”, issued by the Aerial Agricultural Association of Australia Ltd.

### Checklist:

- Have you cleaned/decontaminated your boom sprayer?
- Have you contacted your neighbour prior to spraying?
- Is your sprayer set-up correctly for the particular application?
  - Check
    - boom calibration
    - at nozzle - nozzle choice
    - low drift/what spray quality
    - coarse or larger spray quality?
    - boom height - speed of intended application
    - water volume
- You must check, determine and record the weather conditions immediately prior to, and immediately after the spray application is made.
- Record
  - Temperatures
  - Relative Humidity
  - Delta T
  - Wind speed
  - Is there a temperature inversion?

•Night Spraying - Extra care is required to ensure that inversion conditions are not present. Use smoke generator to determine wind direction and presence of inversion conditions.

For further information refer to [nufarm.com.au/spraywise](http://nufarm.com.au/spraywise)



[spraywisedecisions.com.au](http://spraywisedecisions.com.au) is an online weather forecasting program and is recommended for use when planning your pesticide application.

### Application

Nufarm weedmaster DST is a non-selective translocated herbicide. Direct spray contact, or even slight drift, may cause severe injury or other desirable plants including trees. Clean all equipment after use by thoroughly washing with water.

#### Boom Equipment

For Broadcast boom application, a spray volume of 80 L/ha is recommended for broadcast booms and 200 L/ha or less for treeline and veline spraying in orchards and vineyards. Nozzles and pressure settings should be selected to deliver a COARSE to VERY COARSE spray quality at the target. The use of nozzles and/or pressure settings that produce VERY FINE or FINE spray quality should be avoided as these are prone to loss or drift. In multiple product tank mixes a minimum water volume of 50 L/ha is recommended and local advice should be sought. Correct mixing order is important as is good in-tank agitation when application is occurring.

#### Shielded Equipment

For shielded applications a spray volume of 80 L/sprayed ha is recommended using nozzle types and pressure settings to deliver a COARSE spray quality at the target. Crop damage may result if spray drift occurs through incorrect nozzle and/or pressure selection, inadequate shielding and/or wind strength, high evaporation rates or excessive ground speed.

#### High Volume Application (e.g. Knapsack, Handgun Equipment)

The dilution rate varies depending on the use situation and weeds controlled - see Weeds Controlled tables for specific rates and use recommendation. Adjust equipment to achieve an even spray pattern with a COARSE spray quality at the target. Apply to ensure complete and uniform wetting of all foliage.

#### Controlled Droplet Application Equipment (CDA)

Nufarm weedmaster DST can be applied through hand held and machine mounted CDA sprayers. See Weeds Controlled tables for specific rates and use recommendations. Due to the range of CDA equipment available, dilution rates, flow rates and travel speeds will need to be determined for individual sprayers to ensure labelled rates are applied. Spray units need to be cleaned thoroughly preferably after each application to ensure optimum performance.

**DO NOT** add oils to Nufarm weedmaster DST/water mixture, otherwise difficulty in application and reduced weed control may occur.

Because CDA units may deliver relatively low spray volumes per hectare, use on large weeds may result in insufficient coverage resulting in inadequate weed control.

**CAUTION:** CDA equipment produces a fine spray pattern which is not easily visible. Ensure spray pattern or drift does not contact foliage or any other green tissue of desirable plants, since severe injury or destruction may result.

#### Aerial Equipment

Nufarm weedmaster DST may be applied by aircraft for control of weeds in forests, cropland or pasture prior to establishment of crops, new pastures or new forest plantings and for pre-harvest applications, up to a maximum rate of 3.1 L/ha where specified by this label. **DO NOT** apply treatments by aircraft in situations where drift onto sensitive crops and pastures is likely to occur.

Apply treatments using a spray volume not less than 20 L/ha and using settings to produce a COARSE to VERY COARSE spray quality. In multiple product tank mixes a minimum water volume of 50 L/ha is recommended and local advice should be sought. Correct mixing order is important. Swath width should be set to consider aircraft type, wind conditions and target height. Swath width will need to be reduced to avoid striping under light wind conditions and/or application to tall, dense targets e.g. preharvest application, treatments in heavy crop stubble. Thoroughly wash aircraft after each day of spraying to remove herbicide residues.

When applying this product by helicopter in combination with Associate for the control of Blackberry, Coppice and Pine wilding suppression in forestry and other specific situations, a rate of 6.1 L/ha Nufarm weedmaster DST may be applied. Refer Section 11 and the Associate label for specific recommendations and application recommendations.

#### **Aerial application - Hilly terrain**

For aerial application on hilly terrain, increase water volume to 30-80 L/ha and use a COARSE spray quality to optimise spray coverage.

#### **Aerial application - Air temperature and relative humidity**

**DO NOT** apply Nufarm weedmaster DST by aircraft at temperatures above 30°C. Ensure spray output is at least 30 L/ha when temperatures rise above 25°C. Avoid application when relative humidity falls below 35%.

### **RESISTANT WEEDS WARNING**

## **GROUP 9 HERBICIDE**

Nufarm weedmaster DST Advanced Technology Herbicide is a member of the Glycines group of herbicides. Nufarm weedmaster DST has the inhibition of EPSP synthase mode of action. For weed resistance management Nufarm weedmaster DST is a Group 9 herbicide. Some naturally occurring weed biotypes resistant to Nufarm weedmaster DST and other Group 9 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nufarm weedmaster DST or other Group 9 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm accepts no liability for any losses that may result from the failure of Nufarm weedmaster DST to control resistant weeds.

Users of Nufarm weedmaster DST over Roundup Ready Flex<sup>®</sup> cotton, Roundup Ready<sup>®</sup> canola, TruFlex<sup>®</sup> canola or canola hybrids with the Optimum GLY<sup>®</sup> herbicide tolerance trait must implement practices that minimize the development of resistance in treated weeds. Minimising this risk may best be achieved by following the integrated weed management strategy guidelines summarised below:

1. Aim to enter the Roundup Ready Flex<sup>®</sup> cotton, Roundup Ready<sup>®</sup> canola, TruFlex<sup>®</sup> canola or canola hybrids with the Optimum GLY<sup>®</sup> herbicide tolerance trait, cropping phase of the rotation with a low weed burden.
2. Integrate as many different weed control options (chemical and cultural) as possible through all phases of the crop rotation.
3. Make every herbicide application count – use registered rates at the correct application growth stage and assess effectiveness.
4. Rotate herbicides with different modes of action throughout the crop rotation.
5. Regularly monitor the effectiveness of resistance management practices.
6. Test weed populations for herbicide resistance status as part of ongoing integrated weed management.
7. Growers should not plant Roundup Ready Flex<sup>®</sup> cotton, Roundup Ready<sup>®</sup> canola, TruFlex<sup>®</sup> canola or canola hybrids with the Optimum GLY<sup>®</sup> herbicide tolerance trait in paddocks with populations of confirmed glyphosate resistant weeds.

It is advised that consultation on Integrated Weed Management be undertaken with an accredited agronomist or program prior to use of weedmaster DST over Roundup Ready Flex<sup>®</sup> cotton, Roundup Ready<sup>®</sup> canola, TruFlex<sup>®</sup> canola or canola hybrids with the Optimum GLY<sup>®</sup> herbicide tolerance trait.

More information on Integrated Weed Management can be found at:

- Australian Glyphosate Sustainability Working Group (AGSWG), 'Sustainable glyphosate use in winter grain cropping' <http://glyphosateresistance.org.au>
- Weedsmart: [www.weedsmart.org.au](http://www.weedsmart.org.au)
- CropLife Australia: <http://www.croplife.org.au/industry-stewardship/resistancemanagement/>

As with conventional varieties, volunteer and ratoon Roundup Ready Flex<sup>®</sup> cotton, Roundup Ready<sup>®</sup> canola, TruFlex<sup>®</sup> canola or volunteer canola hybrids with the Optimum GLY<sup>®</sup> herbicide tolerance trait may occur in fallows, and non-cropping areas of a farm such as irrigation ditches, water storages, etc. These plants will not be controlled by Nufarm weedmaster DST or other glyphosate (Group 9) herbicides and should be controlled in both cropping and non-cropping areas. These plants are best managed with cultivation and/or appropriate registered herbicides (see Integrated Weed Management Strategy Guidelines above). Growers should ensure that they have an effective weed management strategy developed for the control of these weeds. Herbicide control options for these plants include the following (refer to product labels for further information on use situations): Agritome<sup>®</sup> 750, Alliance<sup>®</sup>, Amicide<sup>®</sup> Advance 700, Amtrile<sup>®</sup> T, Associate<sup>®</sup>, Bentley<sup>®</sup>, Biffo<sup>®</sup>, Broadside<sup>®</sup>, Broadsword<sup>®</sup>, Bromicide<sup>®</sup> 200, Bromicide<sup>®</sup> MA, Comet<sup>®</sup> 400, Eliminar<sup>®</sup> C, Esterice<sup>®</sup> Xtra 680, Nail<sup>®</sup> 240, Nail<sup>®</sup> 600, Polo<sup>®</sup> 570 LVE, Revolver<sup>®</sup>, Sentry<sup>®</sup>, Shirquat<sup>®</sup> 250, T-Rex<sup>®</sup> and Unity<sup>®</sup>.

#### **Resistant weeds reporting, auditing and surveying**

Users of Nufarm weedmaster DST are required to report any adverse events, such as suspected weed resistance, to Nufarm, as soon as it is identified.

Nufarm will investigate the incident and produce a report of any incidents of confirmed resistance of weeds to Nufarm weedmaster DST in target weed species which are normally susceptible to this herbicide and forward the report as such as practicable to the Australian Pesticides and Veterinary Medicines Authority. Weeds identified to have survived Nufarm weedmaster DST must be controlled by an alternative strategy in order to prevent weeds from setting seed.

Users of Nufarm weedmaster DST over Roundup Ready Flex<sup>®</sup> cotton, Roundup Ready<sup>®</sup> canola, TruFlex<sup>®</sup> canola or canola hybrids with the Optimum GLY<sup>®</sup> herbicide tolerance trait must allow Nufarm or its agents to undertake audits or surveys as necessary to assess management by users of the development of glyphosate resistance in target weeds. Nufarm or its agents will conduct an audit or survey annually on a percentage of fields where Nufarm weedmaster DST has been used over Roundup Ready Flex<sup>®</sup> cotton, Roundup Ready<sup>®</sup> canola, TruFlex<sup>®</sup> canola or canola hybrids with the Optimum GLY<sup>®</sup> herbicide tolerance trait.

### **PRECAUTION**

#### **Re-entry period**

Re-entry is not allowed until the product has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

#### **PROTECTION OF CROPS, NATIVE AND OTHER NONTARGET PLANTS**

Avoid contact with foliage, green stems or fruit of crops, desirable plants and trees, since severe injury or destruction may result. **DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. **DO NOT** apply if wind is blowing towards a sensitive crop or situation and off-target damage cannot be avoided.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

**DO NOT** contaminate dams, rivers or streams with the product or used container. **DO NOT** apply to weeds growing in or over water. **DO NOT** spray across open bodies of water.

#### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

##### For non-returnable containers:

Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.

##### For RETURNABLE containers:

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

#### SAFETY DIRECTIONS

Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles, and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

#### ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

**Causes serious eye irritation. Causes skin irritation.** Wash hands and exposed skin thoroughly after handling. Wear protective gloves, clothing, eye and face protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before reuse.

#### SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained by scanning the QR code, from your supplier or the Nufarm website.

**In case of emergency: Phone 1800 033 498 (24 hrs) and ask for shift supervisor.**

#### PRODUCT STEWARDSHIP INFORMATION AND TOOLS

[nufarm.com.au/spraywise](http://nufarm.com.au/spraywise) is a website which contains resources on spray drift management, recording keeping and application technologies.  
[spraywisedecisions.com.au](http://spraywisedecisions.com.au) is an online weather forecasting program and is recommended for use when planning your pesticide application.  
When spraying in or near areas with sensitive crops such as cotton, check online at [satacrop.com.au](http://satacrop.com.au) for the proximity of sensitive crops mapped by growers.  
[stewardshipfirst.com.au](http://stewardshipfirst.com.au) is a website with CroLife Australia's suite of world-leading product stewardship initiatives, programs and best-practice guides.

#### CONDITIONS OF SALE

Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.

\* All trade marks owned or used under licence by Nufarm Australia Limited

\* Other registered trade marks



**AUSTRALIAN  
THROUGH  
& THROUGH**

**nufarm.com.au**

**NUFARM AUSTRALIA LIMITED**

103-105 Pipe Road, Laverton North Victoria 3026

For product advice, or to learn more about complementary Nufarm solutions,  
reach out to your local Nufarm specialist or call **1800 NUFARM (1800 683 267)**.

APVMA Approval No.: 62723/133612

Date: 07-02-2022