

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Talon[®] GT Pro

Rodenticide Grain Bait Block

syngenta[®]

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM

Ready to use block bait for use in and around industrial, commercial, public services, agricultural and domestic buildings

Controls rodents resistant to other anticoagulants, including warfarin, coumatetralyl and bromadiolone

ONE FEED KILLS RATS AND MICE

**Red Grain
BLOCKS**



Syngenta Australia Pty Ltd

Level 1, 2 Lyonpark Road, Macquarie Park NSW 2113

In a transport emergency dial 000, Police or Fire Brigade

**For specialist advice in an emergency only, call 1800 033 111
(24 hours)**

APVMA Approval No: 92319/135173

®

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply bait to crops

DO NOT place bait in the open

DO NOT exceed 3 m between bait stations for mice, or 9 m for rats.

Baiting Strategy

Eliminate as far as practicable all alternative food sources. Bait infested area. Place bait stations under cover. Inspect bait stations regularly. Replace eaten baits. If eaten quickly, increase the number of bait stations. Continue observation and replenishment until no more bait is taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Situation	Pest	Rate	Critical Comments
Agricultural Commercial Domestic Industrial Buildings	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>) Mice (<i>Mus musculus</i>)	1 to 3 blocks per bait station depending on level of infestation	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings. For rats: Place 1-3 blocks at intervals of 5-9 m in infested areas. For mice: Place single blocks at 2-3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Ensure baits are inaccessible to children, pets, domestic animals and wildlife. Place bait along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if appropriate. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and location of bait locations for the purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait. Regularly search for, remove and destroy dead rodents, by burning or burying them. DO NOT place in refuse bins or rubbish tips. Ensure bait blocks are removed when rodent activity has ceased.

PRECAUTION

DO NOT place bait in areas where there is a possibility of directly contaminating food, food utensils, livestock feed or surfaces that may come in direct contact with food.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Depending on the amount eaten, this bait may kill or injure. The bait and dead or affected rodents are hazardous to dogs, cats, pigs, poultry and other wildlife. DO NOT place baits in locations which are accessible to pets, domestic animals, livestock, birds or other non-target wildlife. DO NOT contaminate streams, rivers and waterways with the bait or its used container. The use of this product for the control of protected native rodents and rodent like animals requires permission from wildlife authorities. The product is restricted to use in and around buildings (within 2 m) or enclosed spaces, eg drains.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food and feedstuffs, seed and fertilisers.

Containers which have held bait should not be used for any other purpose. Triple-rinse containers before disposal and rinsate should be disposed of in the disposal pit specifically marked and set up for this purpose, clear of water ways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Dispose of empty containers and unused or untaken bait by delivering to an approved waste management facility. If an approved waste management facility is not available, bury the unused or untaken bait and empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. Search for dead rats or mice – burn or bury them. DO NOT place in refuse bins or rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when rodent activity has ceased.

SAFETY DIRECTIONS

Poisonous if swallowed. Do not inhale dust. DO NOT touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K1 [phytomenadione] is antidotal.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126; Vitamin K1 (Phytomenadione) is antidotal.

NOTE TO PHYSICIANS AND VETERINARIANS

Brodifacoum is a long-lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate paramedical personnel or veterinarians. If administered intravenously the injection must be given slowly.

SAFETY DATA SHEET

If additional hazard information is required refer to the Safety Data Sheet. For a copy visit our website at www.syngentappm.com.au or www.syngenta.com.au or use the QR code on this label.

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



Batch No.:

Date of Manufacture: