

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Advion[®] Evolution

Cockroach Gel

syngenta[®]

ACTIVE CONSTITUENT: 8 g/kg INDOXACARB (75:25)
(equivalent to 6 g/kg active S-isomer)

GROUP	22A	INSECTICIDE
-------	------------	-------------

For the control of Cockroaches in the situations specified
in the Directions for Use

FOR PROFESSIONAL USE ONLY

Syngenta Australia Pty Ltd
Level 1, 2 Lyonpark Road, Macquarie Park NSW 2113

In a transport emergency dial 000, Police or Fire Brigade
For specialist advice in an emergency only, call 1800 033 111 (24 hours)

APVMA Approval No: 91623/132864

®

DIRECTIONS FOR USE

Restraints

DO NOT place gel on surfaces that are regularly washed

DO NOT allow to contact food, livestock feed or their water

DO NOT place gel on surfaces where food is prepared or on utensils that are used for food preparation, on food packing materials or on surfaces where food packing materials are stored

DO NOT apply ADVION® Evolution Cockroach Gel to surfaces where the temperature exceeds 50°C as this could cause the gel to liquefy and lose effectiveness. (Examples include, but are not limited to, parts of stoves, ovens, grills, fume hoods, saunas, heat lamps, coffee urns, toasters, fryers, dishwashers and hot water pipes)

Situation	Pest	Rate	Critical Comments
<p>Domestic, commercial and public service buildings including factories, farm buildings, food processing establishments, homes, hospitals, kitchens, offices, store rooms, restaurants, retail outlets</p> <p>Vehicles, Other situations which fall within these definitions</p>	<p>American Cockroach (<i>Periplaneta americana</i>), German Cockroach (<i>Blattella germanica</i>), other common cockroaches</p>	<p>Apply at least 2 spots/m². Increasing the number of spots (up to 5 spots/m²) may be required for heavy infestations or if American Cockroaches are the main pest. Recommended spot size is approximately 5 mm diameter (0.1 g)</p>	<p>Infested areas should be carefully inspected to determine overall infestation levels, locations of harbourage areas and most appropriate application points. ADVION® Evolution Cockroach Gel should be applied as a small spot of gel, approximately 0.1 g (approximately 5 mm in diameter – the size of a small pea), to areas not able to be contacted by children or pets. Several small applications distributed within an area are more effective than 1 or 2 applications of large quantities. In some cases a small thin bead (less than 2.5 mm wide and 5 cm long) may be appropriate, such as between construction elements or other cracks and crevices. Indoor applications can include cracks and crevices along walls or floors, behind or under equipment, under tables, within framing of tables or furniture, around sinks, within or near garbage collection areas, attics, crawl space and cracks and crevices in cabinets. Bait placements should be inspected periodically for consumption or attractiveness and reapplied as needed. A typical infestation may require as little as 15 g of Advion Evolution Cockroach Gel whereas heavy infestations in large structures may require the use of more of the gel bait.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

GENERAL INSTRUCTIONS

ADVION® Evolution Cockroach Gel is designed to control pest species of cockroaches including American and German. The slight delay in mortality caused by the active ingredient, indoxacarb, allows cockroaches to consume the bait and return to the harbourage site to contaminate other cockroaches resulting in significant reduction in population levels.

Use Instructions and Precautions

- ADVION® Evolution Cockroach Gel should be applied to areas frequented by cockroaches such as between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, and behind and under appliances, cabinets, sinks, closets and any other equipment that cockroaches are typically known to hide.
- Avoid applying ADVION® Evolution Cockroach Gel to surfaces recently treated with repellent residual sprays.
- Surfaces treated with ADVION® Evolution Cockroach Gel should not be treated with repellent residual sprays.
- Repeat application may be made after follow up inspections of the treated areas.
- DO NOT apply to surfaces that food contacts.
- DO NOT allow open foods to contact gel material.

To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated and slowly depress plunger until sufficient gel is dispensed. Recap the dispenser after treatment is completed.

Where maintenance treatments form part of the professional pest controller's contract, it is recommended that cockroach activity be monitored where gel had been applied 1 month after initial treatment or at more regular intervals in areas of high infestation.

Insecticide Resistance Warning

GROUP	22A	INSECTICIDE
--------------	------------	--------------------

For insecticide resistance management ADVION® Evolution Cockroach Gel is a Group 22A insecticide. Some naturally occurring insect biotypes resistant to ADVION® Evolution Cockroach Gel and other Group 22A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if ADVION® Evolution Cockroach Gel and other Group 22A insecticides are used repeatedly. The effectiveness of ADVION® Evolution Cockroach Gel on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from failure of ADVION® Evolution Cockroach Gel to control resistant insects. ADVION® Evolution Cockroach Gel may be subject to specific resistance management strategies. For further information contact your supplier, Syngenta representative or local government entomologist.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

DO NOT apply this product in any areas easily accessible to children or animals.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated locked place out of the reach of children. DO NOT store in direct sunlight and DO NOT store at temperatures exceeding 35°C. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in the garbage.

SAFETY DIRECTIONS

Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

SAFETY DATA SHEET

If additional hazard information is required refer to the Safety Data Sheet. For a copy visit our website at www.syngentappm.com.au or www.syngenta.com.au or use the QR code on this label.

May cause skin sensitisation by skin contact
--

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

