

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING

Active Constituent : 750 g/kg HALOSULFURON-METHYL

GROUP	B	HERBICIDE
-------	----------	-----------

For selective post-emergence control of Nutgrass in Sugarcane, Corn/Maize and Sorghum and for the selective post-emergence control of Nutgrass and Mullumbimby couch in Turf. Also for the control of Nutgrass in Cotton using shielded sprayers.



Distributed by
Nufarm Australia Limited ACN 004 377 780
103-105 Pipe Road, Laverton North, VIC 3026
Tel: (03) 9282 1000 Fax: (03) 9282 1001
NRA Approval No 56620/0603

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Dust will irritate the eyes. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre Tel 13 11 26.

<p>ONLY IN EMERGENCY CALL 1 800 033 498 ALL HOURS Refer to Material Safety Data Sheet</p>

For Technical enquiries
phone 1800 639 899

Batch number:

Date of manufacture :

NRA Approval No 56620/0603

CONDITION OF SALE

Nufarm Australia Limited ("Nufarm") shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Nufarm's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to alter these conditions.

SUGARCANE (NSW AND QLD ONLY)

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Plant and ratoon Sugarcane	Nutgrass	65-130 g/ha	Use the higher rate for dense infestations or for maximum control where a single application is intended. Follow-up treatments may be required to control plants emerging from dormant tubers. Apply using a boom spray with flat fan nozzles or flood jets to apply at least 80 L/ha as a broadcast, or directed treatment in Sugarcane. Avoid streaking, skips or overlaps during application. Cultivation should be delayed for at least 2 days following treatment.
		1.3 g/100sq m	For spot treatment using handgun or knapsack sprayers, apply 1.3g Sempra per 100 square meters. For example, mix 1.3 g Sempra in 10L of water and apply 10L of the mix per 100 square meters.

CORN/MAIZE (ZEA MAYS, ALL STATES, ACT, NT)

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Maize/Corn	Nutgrass	65-130 g/ha	<p>Use the higher rate for dense infestations or for maximum control where a single application is intended. Apply post emergence to crop at any time up to a crop height of 60cm. Apply using a boom spray with flat fan nozzles or flood jets to apply at least 80 L/ha as a broadcast or directed treatment. Time treatment to ensure maximum emergence of Nutgrass when majority of plants are 3-4 leaf stage. Some transient yellowing and growth reduction of crop may occur at higher rate.</p> <p>Do not apply Sempra to sweetcorn, popcorn or hybrids. Do not apply Sempra herbicide to corn/maize if the crop is under severe stress due to drought, water saturated soil, low fertility (especially low nitrogen levels) or other poor growing conditions.</p>

SORGHUM (NSW AND QLD ONLY)

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Sorghum	Nutgrass	65-130 g/ha	Use the higher rate for dense infestations or for maximum control where a single application is intended. Apply using a boom spray with flat fan nozzles or flood jets to apply at least 80 L/ha as a broadcast or directed treatment. Time treatment to ensure maximum emergence of Nutgrass when majority of plants are 3-4 leaf stage. Apply post emergence to crop at any time up to a crop height of 60cm. Do not apply more than 200 g/ha per season. Apply as a directed spray if using rates greater than 65 g/ha. Some transient yellowing and growth reduction of crop may occur at higher rate.

COTTON (NSW AND QLD ONLY)

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Shielded sprayer application in irrigated cotton only.	Nutgrass	65-130 g/ha	Use the higher rate for dense infestations or for maximum control where a single application is intended. Apply Sempra to Nutgrass growing between Cotton rows using a shielded sprayer. Apply in crops at least 20cm high but before first flower. Spray or spray drift which contacts any part of the Cotton plant may cause severe injury. Follow-up treatments may be required to control plants emerging from dormant tubers. Cultivation should be delayed for at least 2 days following treatment. The first irrigation after spraying should be delayed for as long as possible to minimise herbicide runoff.
Shielded or directed handgun or knapsack application		1.3 g/100sq m	For spot treatment using shielded or directed handguns or knapsack sprayers. DO NOT apply in crops less than 20cm high. Spray or spray drift which contacts any part of the Cotton plant may cause severe injury. Apply 1.3 g Sempra per 100 square meters. For example, mix 1.3 g Sempra in 10L of water and apply 10L of the mix per 100 square meters.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL, UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

HARVEST

TURF: DO NOT USE CLIPPINGS FROM TREATED AREAS FOR MULCHING AROUND VEGETABLES OR FRUIT TREES. DO NOT FEED GRASS CLIPPINGS FROM TREATED AREAS TO POULTRY OR OTHER LIVESTOCK.

ALL OTHER CROPS: WITHHOLDING PERIOD NOT REQUIRED WHEN IS USED AS DIRECTED.

GRAZING

SORGHUM: DO NOT GRAZE LIVESTOCK OR CUT FOR FODDER OF FORAGE FOR FOUR WEEKS AFTER TREATMENT.

TURF: DO NOT ALLOW GRAZING OF TREATED TURF.

ALL OTHER CROPS: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Sempra is a dry flowable granule which disperses in water. Sempra can be used for selective post-emergence control of Nutgrass in Sugarcane, Corn/Maize and Sorghum and for the selective post emergence control of Nutgrass and Mullumbimby couch in Turf. Sempra may also be used for the control of Nutgrass in Cotton using shielded sprayers.

Symptoms of weed control are a gradual yellowing of foliage and seed heads followed by desiccation. Initial symptoms may take 7-10 days to be noticeable, with full effects occurring 4 to 6 weeks after treatment. Sempra should be applied to actively growing weeds when new growth has reached a minimum of 5cm of new leaf growth for Nutgrass or 2cm for Mullumbimby couch. Apply follow-up treatments if sufficient new growth warrants re-treatment. Irrigation or rainfall within two hours of application will reduce control. Drought stress after treatment may also reduce control.

RESISTANT WEEDS WARNING

Sempra Herbicide inhibits acetolactate synthase and therefore is a member of the sulfonylurea group of herbicides. When considering resistant weed management, Sempra Herbicide is a Group B herbicide. Some naturally occurring weed biotypes resistant to Sempra Herbicide and other inhibitors of acetolactate synthase may exist through normal genetic variability in any weed population. Resistant weeds can eventually dominate the weed population if group B herbicides are used repeatedly and as a consequence, resistant weeds will not be controlled by Sempra Herbicide or any other herbicides that inhibit acetolactate synthase. Since development of resistant weeds is difficult to detect prior to herbicide use, Nufarm accepts no liability for any losses that may result from the failure of Sempra Herbicide to control resistant weeds.

MIXING

Sempra is a dry flowable granule that disperses in water. Add the measured amount gradually to a part-filled spray tank while maintaining continuous bypass agitation. Add the surfactant near the end of the filling process to avoid excessive foaming. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the water source. If allowed to stand, ensure that the mixture is thoroughly agitated before re-commencing spraying. Use the mixture within one day.

SPRAYER CLEANUP

Before application of products other than Sempra the sprayer must be cleaned out as follows:

1. Drain the tank and flush equipment with water for a minimum of 10 minutes, including hoses, filters and booms.
2. Fill the tank with clean water and add chlorine bleach (contains 4% chlorine) at the rate of 300mL/100L of water. Flush through the boom and agitate for 15 minutes.
3. Repeat step 2 above.
4. Remove all nozzles and screens and clean thoroughly.
5. To remove traces of chlorine bleach, rinse the tank thoroughly with clean water and flush through hoses and booms.

Caution: Do not use chlorine bleach with ammonia. All traces of liquid fertilizer containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odour, which can cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

SURFACTANT ADDITION

Sempra must be applied with a non-ionic surfactant to ensure uptake. Use 200mL/100L of a 600g/L non-ionic surfactant or equivalent. For hand-gun or knapsack application, add surfactant at 20mL/10L of water.

COMPATIBILITY

Sempra is compatible with Roundup®, Roundup® CT, Roundup PowerMAX™ and Roundup® Dry, Hoegrass, dicamba/MCPA and bromoxynil/MCPA mixtures and amine formulations of 2,4-D. Read and follow all label directions, restraints, plant-back periods, withholding periods, regional use restrictions and safety directions for the tank mix products. Tank mixing with paraquat or ametryn may result in reduced weed control. Organophosphate insecticides should not be tank-mixed or applied 7 days prior or 3 days after application of Sempra, as severe crop injury may result.

FOLLOWING CROPS

The following crops may be planted at specific time intervals following application of approved rates of Sempra in approved situations. Use the time intervals listed below to determine the required time interval before planting.

Crop	Plant back interval after the last application of Sempra
Corn/Maize, Sorghum and Sugarcane	2 months
Wheat	3 months
Cotton	4 months
All other crops (except sugarbeet)	24 months
Sugarbeet	36 months

PROTECTION OF CROP, NATIVE AND OTHER NON-TARGET PLANTS

Avoid spraying of non-target vegetation. In turf, do not use clippings from treated areas for mulching around vegetables or fruit trees. DO NOT apply under weather conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. After spraying, the first flush irrigation tailwater or stormwater run-off from land treated with Sempra should be prevented from entering waterways. Where farms do not have the capacity to retain run-off, DO NOT irrigate to the point of run-off for at least 6 days after application.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEA, THE ENVIRONMENT AND OTHERS

Do not contaminate dams, rivers or streams with the product or used container. Do not feed grass clippings from treated areas to poultry or other livestock or allow grazing of treated turf. Keep people and pets off treated areas until the spray solution has dried. Sempra is very highly toxic to duckweed.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Dispose of in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Dust will irritate the eyes. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre Tel 13 11 26.

<p>ONLY IN EMERGENCY CALL 1 800 033 498 ALL HOURS Refer to Material Safety Data Sheet</p>

For Technical enquiries
phone 1800 639 899

NRA Approval No 56620/ 0603

CONDITION OF SALE

Nufarm Australia Limited ("Nufarm") shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Nufarm's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to alter these conditions.

Roundup and Roundup PowerMAX are registered / trademark of Monsanto Technology LLC used under licence by Nufarm Australia Limited., Sempra is a registered trademark of Nissan Chemical Industries Ltd. Copyright Nufarm Australia Limited, May 2003